

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 535 910 A1

(12)

EUROPEAN PATENT APPLICATION

published in accordance with Art. 158(3) EPC

(43) Date of publication:

01.06.2005 Bulletin 2005/22

(21) Application number: **03721022.6**

(22) Date of filing: **01.05.2003**

(51) Int Cl.7: **C07D 215/22**, C07D 417/12,
A61K 31/47, A61K 31/4709,
A61K 31/5377, A61P 19/02,
A61P 19/10, A61P 29/00,
A61P 35/00, A61P 35/04,
A61P 43/00

(86) International application number:

PCT/JP2003/005593

(87) International publication number:

WO 2003/093238 (13.11.2003 Gazette 2003/46)

(84) Designated Contracting States:

**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IT LI LU MC NL PT RO SE SI SK TR**

Designated Extension States:

AL LT LV MK

(30) Priority: **01.05.2002 JP 2002130049**

(71) Applicant: **KIRIN BEER KABUSHIKI KAISHA**
Tokyo 104-8288 (JP)

(72) Inventors:


- **KUBO, Kazuo,**
Pharmaceutical Research Laboratories
Takasaki-shi, Gunma 370-1295 (JP)

- **OHNO, Hiroaki,**
Pharmaceutical Research Laborator.
Takasaki-shi, Gunma 370-1295 (JP)
- **ISOE, Toshiyuki,**
c/o Kirin Beer Kabushiki Kaisha
Tokyo 150-8011 (JP)
- **NISHITOBA, Tuyoshi,**
Kirin Beer Kabushiki Kaisha
Tokyo 150-8011 (JP)

(74) Representative: **HOFFMANN EITLE**
Patent- und Rechtsanwälte
Arabellastrasse 4
81925 München (DE)

(54) **QUINOLINE DERIVATIVES AND QUINAZOLINE DERIVATIVES INHIBITING
AUTOPHOSPHORYLATION OF MACROPHAGE COLONY STIMULATING FACTOR RECEPTOR**

(57) An objective of the present invention is to provide compounds which have inhibitory activity against autophosphorylation of macrophage colony-stimulating factor receptors. The compounds of the present invention are represented by formula (I) and salt and solvate thereof:



(I)

EP 1 535 910 A1

wherein X represents CH or N; Z represents O or S; R¹, R², and R³ represent H, optionally substituted alkoxy or the like; R⁴ represents H; R⁵, R⁶, R⁷, and R⁸ represent H, halogen, alkyl, alkoxy, trifluoromethyl or the like; R⁹ and R¹⁰ represent H, alkyl or the like; and any one of R¹¹ and R¹² represents H with the other representing alkyl and R¹³ represents an optionally substituted carbocyclic or heterocyclic ring or the like, or R¹¹ represents H and R¹² and R¹³ combine together to form a bicyclic carbocyclic ring.

Description

BACKGROUND OF THE INVENTION

Field of the Invention

[0001] The present invention relates to quinoline derivatives and quinazoline derivatives having inhibitory activity against macrophage colony-stimulating factor receptor autophosphorylation, and more particularly to quinoline derivatives and quinazoline derivatives that are useful for the treatment and prevention of diseases such as bone metastasis of malignant tumors including breast cancer, prostatic cancer, and lung cancer, multiple myeloma, osteoporosis, Behcet's disease, rheumatoid arthritis or other diseases.

Related Art


[0002] Quinoline derivatives and quinazoline derivatives are disclosed in WO 97/17329, Japanese Patent Application No. 328782/1997, and WO 00/43366.

SUMMARY OF THE INVENTION

[0003] The present inventors have found that a group of quinoline derivatives and quinazoline derivatives have high inhibitory activity against macrophage colony-stimulating factor receptor autophosphorylation. Macrophage colony-stimulating factor receptor autophosphorylation is known to be involved, e.g., in the onset of diseases such as bone metastasis of malignant tumors including breast cancer, prostatic cancer, and lung cancer; multiple myeloma; osteoporosis; Behcet's disease; and rheumatoid arthritis.

[0004] An object of the present invention is to provide a compound having inhibitory activity against macrophage colony-stimulating factor receptor autophosphorylation.

[0005] According to the present invention, there are provided compounds represented by formula (I) or pharmaceutically acceptable salts or solvates thereof:



wherein

X represents CH or N;

Z represents O or S;

R¹, R², and R³, which may be the same or different, represent a hydrogen atom; a halogen atom; hydroxyl; cyano; C₁₋₆ alkyl; C₁₋₆ alkoxy; C₂₋₆ alkenyl; C₂₋₆ alkynyl; nitro; -NR¹⁰⁶R¹⁰⁷ wherein R¹⁰⁶ and R¹⁰⁷, which may be the same or different, represent a hydrogen atom or C₁₋₄ alkyl in which the alkyl group is optionally substituted by hydroxyl, -OR¹⁰⁸ wherein R¹⁰⁸ represents C₁₋₄ alkyl, or -NR¹⁰⁹R¹¹⁰ wherein R¹⁰⁹ and R¹¹⁰, which may be the same or different, represent a hydrogen atom or C₁₋₄ alkyl; -CONR¹¹¹R¹¹² wherein R¹¹¹ and R¹¹², which may be the same or different, represent a hydrogen atom or C₁₋₄ alkyl in which the alkyl group is optionally substituted by hydroxyl, -OR¹¹³ wherein R¹¹³ represents C₁₋₄ alkyl, or -NR¹¹⁴R¹¹⁵ wherein R¹¹⁴ and R¹¹⁵, which may be the same or different, represent a hydrogen atom or C₁₋₄ alkyl; or -COOR¹¹⁶ wherein R¹¹⁶ represents a hydrogen atom or C₁₋₄ alkyl in which the alkyl group is

optionally substituted by hydroxyl, $-OR^{117}$ wherein R^{117} represents C_{1-4} alkyl, or $-NR^{118}R^{119}$ wherein R^{118} and R^{119} , which may be the same or different, represent a hydrogen atom or C_{1-4} alkyl in which the C_{1-6} alkyl, C_{1-6} alkoxy, C_{2-6} alkenyl, and C_{2-6} alkynyl groups are optionally substituted by a halogen atom; hydroxyl; C_{1-4} alkyl; C_{1-4} alkoxy; C_{1-4} alkoxy carbonyl; amino in which one or two hydrogen atoms on the amino group each are optionally substituted by C_{1-4} alkyl optionally substituted by hydroxyl or C_{1-4} alkoxy; group $R^{15}R^{16}N-C(=O)-O-$ wherein R^{15} and R^{16} , which may be the same or different, represent a hydrogen atom or C_{1-4} alkyl in which the alkyl group is optionally substituted by hydroxyl or C_{1-4} alkoxy; or group $R^{17}-(S)_m-$ wherein R^{17} represents a saturated or unsaturated three- to seven-membered carbocyclic or heterocyclic group optionally substituted by a halogen atom or C_{1-4} alkyl and m is 0 (zero) or 1,

R^4 represents a hydrogen atom,

R^5 , R^6 , R^7 , and R^8 , which may be the same or different, represent a hydrogen atom, a halogen atom, C_{1-4} alkyl, C_{1-4} alkoxy, C_{1-4} alkylthio, trifluoromethyl, nitro, or amino,

R^9 and R^{10} , which may be the same or different, represent a hydrogen atom, C_{1-6} alkyl, or C_{1-4} alkyl carbonyl, and any one of R^{11} and R^{12} represents a hydrogen atom while the other represents C_{1-4} alkyl, and R^{13} represents a

saturated or unsaturated three- to seven-membered carbocyclic or heterocyclic group or a saturated or unsaturated nine- to twelve-membered bicyclic carbocyclic group in which the carbocyclic and heterocyclic groups are optionally substituted by a halogen atom; hydroxyl; C_{1-4} alkyl; C_{1-4} alkoxy; C_{1-4} alkylthio; trifluoromethyl; nitro; or $-NR^{137}R^{138}$ wherein R^{137} and R^{138} , which may be the same or different, represent a hydrogen atom or C_{1-4} alkyl in which the alkyl group is optionally substituted by hydroxyl, $-OR^{139}$ wherein R^{139} represents C_{1-4} alkyl, or $-NR^{140}R^{141}$ wherein R^{140} and R^{141} , which may be the same or different, represent a hydrogen atom or C_{1-4} alkyl, or

R^{11} represents a hydrogen atom, and R^{12} and R^{13} may combine with a carbon atom attached thereto to form a saturated or unsaturated nine- to twelve-membered bicyclic carbocyclic group.

[0006] The compounds according to the present invention are useful for the treatment and prevention of diseases such as bone metastasis of malignant tumors including breast cancer, prostatic cancer, and lung cancer, multiple myeloma, osteoporosis, Behcet's disease, rheumatoid arthritis or other diseases.

BRIEF DESCRIPTION OF THE DRAWINGS

[0007]

Fig. 1 is a diagram showing concentration-dependent inhibition of M-CSF-dependent cell growth of BAC-1.2F5 cell strain by compounds of the present invention, wherein closed squares represent data on the compound of Example 17, closed rhombuses represent data on the compound of Example 18, and closed triangles represent the compound of Example 74, and the values are average value \pm SE;

Fig. 2 is a diagram showing concentration-dependent inhibition of osteoclast differentiation by the compound of the present invention (Example 74), wherein the values are average value \pm SE;

Fig. 3 is a diagram showing the effect of inhibiting the expansion of bone resorption area involved in bone metastasis of melanoma of compounds according to the present invention in a nude mouse bone metastasis model, wherein closed squares represent data on the compound of Example 74 (20 mg, number of doses: twice), and closed rhombus data on a medium; the values are average value \pm SE; and **: $p < 0.01$ (vs. vehicle control);

Fig. 4 is a diagram showing the effect of inhibiting the expansion of bone resorption area involved in bone metastasis of melanoma of compounds according to the present invention in a nude rat bone metastasis model, wherein the values are average value \pm SE, and *: $p < 0.05$ (vs. vehicle control); and

Fig. 5 is a diagram showing the effect of inhibiting an increase in the number of osteoclasts by a compound of the present invention in an ovariectomy model rats, wherein OVX represents a group of ovariectomy rats and Sham a group of sham operation rats; the values are average value \pm SE; and *: $p < 0.05$ (vs. OVX-vehicle).

DETAILED DESCRIPTION OF THE INVENTION

Compound

[0008] The terms " C_{1-6} alkyl" and " C_{1-6} alkoxy" as used herein as a group or a part of a group respectively mean straight chain or branched chain alkyl and alkoxy having 1 to 6 carbon atoms, preferably 1 to 4 carbon atoms.

[0009] The terms " C_{2-6} alkenyl" and " C_{2-6} alkynyl" as used herein as a group or a part of a group respectively mean straight chain or branched chain alkenyl and alkynyl having 2 to 6 carbon atoms, preferably 2 to 4 carbon atoms.

[0010] Examples of C_{1-6} alkyl include methyl, ethyl, n-propyl, isopropyl, n-butyl, i-butyl, s-butyl, t-butyl, n-pentyl, and n-hexyl.

[0011] Examples of C_{1-6} alkoxy include methoxy, ethoxy, n-propoxy, i-propoxy, n-butoxy, i-butoxy, s-butoxy, and t-butoxy.

[0012] Examples of C₂₋₆ alkenyl include allyl, butenyl, pentenyl, and hexenyl.

[0013] Examples of C₂₋₆ alkynyl include 2-propenyl, butynyl, pentynyl, and hexynyl.

[0014] The term "halogen atom" means a fluorine, chlorine, bromine, or iodine atom.

[0015] The saturated or unsaturated three- to seven-membered carbocyclic or heterocyclic ring is preferably a five- to seven-membered, more preferably five- or six-membered, saturated or unsaturated carbocyclic or heterocyclic ring.

[0016] The saturated or unsaturated nine- to twelve-membered bicyclic carbocyclic or heterocyclic ring is preferably nine- to eleven-membered, more preferably nine- or ten-membered, saturated or unsaturated bicyclic carbocyclic or heterocyclic ring.

[0017] Examples of the saturated or unsaturated three- to seven-membered carbocyclic group include phenyl, cycloheptyl, cyclohexyl, and cyclopentyl.

[0018] Examples of the saturated or unsaturated nine- to twelve-membered bicyclic carbocyclic group include naphthyl, 1,2,3,4-tetrahydronaphthyl, and indanyl.

[0019] The saturated or unsaturated three- to seven-membered heterocyclic ring contains at least one hetero-atom selected from oxygen, nitrogen, and sulfur atoms. The term "hetero-atom" as used herein means oxygen, nitrogen, and sulfur atoms. Examples of the saturated or unsaturated three- to seven-membered heterocyclic group include pyridyl, piperidino, piperazino, morpholino, imidazolyl, triazolyl, tetrazolyl, oxazolyl, thiazolyl, pyrrolidinyl, pyrazolyl, isoxazolyl, and isothiazolyl.

[0020] R¹ preferably represents a hydrogen atom.

[0021] C₁₋₆ alkyl, C₁₋₆ alkoxy, C₂₋₆ alkenyl, and C₂₋₆ alkynyl, which R¹, R², and R³ may represent, are optionally substituted by R¹⁷-(S)_m- wherein m is preferably 0. The carbocyclic group and the heterocyclic group, which R¹⁷ may represent, preferably represent a saturated or unsaturated five- or six-membered carbocyclic or heterocyclic group. More preferably, the carbocyclic group represents phenyl. More preferably, the heterocyclic group represents a saturated or unsaturated five-membered heterocyclic group containing 1 to 4 nitrogen atoms or a saturated or unsaturated six-membered heterocyclic group containing 1 to 2 hetero-atoms selected from nitrogen and oxygen atoms. The hetero-atom constituting the six-membered heterocyclic group may be more specifically one nitrogen atom and one oxygen atom, or one or two nitrogen atoms.

[0022] When m is 0 (zero), -(S)_m- represents a single bond.

[0023] The substituted C₁₋₆ alkoxy, which R¹, R² and R³ may represent, preferably represents group R³¹-(CH₂)_p-O- wherein R³¹ represents a halogen atom; hydroxyl; C₁₋₄ alkyl; C₁₋₄ alkoxy; C₁₋₄ alkoxy carbonyl; amino in which one or more hydrogen atoms on the amino group each are optionally substituted by C₁₋₄ alkyl optionally substituted by hydroxyl or C₁₋₄ alkoxy; group R¹⁵R¹⁶N-C(=O)-O- wherein R¹⁵ and R¹⁶ are as defined in formula (I); or group R¹⁷-(S)_m- wherein R¹⁷ is as defined in formula (I), and p is an integer of 1 to 6, preferably 1 to 4.

[0024] R² and R³, which may be the same or different, preferably represent C₁₋₆ alkoxy. This alkoxy group is optionally substituted by a halogen atom; hydroxyl; C₁₋₄ alkyl; C₁₋₄ alkoxy; C₁₋₄ alkoxy carbonyl; amino in which one or two hydrogen atoms on the amino group each are optionally substituted by C₁₋₄ alkyl optionally substituted by hydroxyl or C₁₋₄ alkoxy; or a saturated or unsaturated three- to seven-membered carbocyclic or heterocyclic group.

[0025] R² and R³, which may be the same or different, preferably represent C₁₋₆ alkoxy optionally substituted by R¹⁷-(S)_m-, more preferably methoxy or R³¹-(CH₂)_p-O-.

[0026] X preferably represents CH, and Z preferably represents O.

[0027] Preferably, at least one of R⁵, R⁶, R⁷, and R⁸ represents a halogen atom, more preferably a chlorine or fluorine atom, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio, trifluoromethyl, nitro, or amino and the remaining groups represent a hydrogen atom.

[0028] Preferably, all of R⁵, R⁶, R⁷, and R⁸ represent a hydrogen atom.

[0029] R⁵ and R⁶, which may be the same or different, preferably represent a halogen atom, more preferably a chlorine or fluorine atom, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio, trifluoromethyl, nitro, or amino, and R⁷ and R⁸ represent a hydrogen atom.

[0030] R⁶ and R⁷, which may be the same or different, preferably represent a halogen atom, more preferably a chlorine or fluorine atom, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio, trifluoromethyl, nitro, or amino, and R⁵ and R⁸ represent a hydrogen atom.


[0031] R⁹ and R¹⁰ preferably represent a hydrogen atom.

[0032] Preferred examples of the carbocyclic group, which R¹³ may represent, include phenyl and naphthyl.

[0033] Preferred examples of the heterocyclic group, which R¹³ may represent, include imidazolyl, oxazolyl, thiazolyl, pyrazolyl, isoxazolyl, and isothiazolyl.

[0034] Examples of the bicyclic carbocyclic group, which R¹² and R¹³ may represent together with a carbon atom attached thereto, include 1,2,3,4-tetrahydronaphthyl and indanyl.

[0035] A group of preferred compounds represented by formula (I) include a group of compounds represented by formula (Ia):



wherein

X represents CH or N,

R¹⁸ and R¹⁹, which may be the same or different, represent C₁₋₆ alkoxy, said alkoxy group being optionally substituted by a halogen atom; hydroxyl; C₁₋₄ alkyl; C₁₋₄ alkoxy carbonyl; amino in which one or two hydrogen atoms on the amino group each are optionally substituted by C₁₋₄ alkyl optionally substituted by hydroxyl or C₁₋₄ alkoxy; or a saturated or unsaturated three- to seven-membered carbocyclic or heterocyclic group,

R²⁰, R²¹, R²², and R²³, which may be the same or different, represent a hydrogen atom, a halogen atom, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio, trifluoromethyl, nitro, or amino,

any one of R²⁴ and R²⁵ represents a hydrogen atom and the other represents C₁₋₄ alkyl, and R²⁶ represents phenyl, naphthyl, imidazolyl, oxazolyl, thiazolyl, pyrazolyl, isoxazolyl, or isothiazolyl, said groups being optionally substituted by a halogen atom, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio, trifluoromethyl, nitro, or amino in which one or two hydrogen atoms on the amino group each are optionally substituted by C₁₋₄ alkyl, or

R²⁴ represents a hydrogen atom, and R²⁵ and R²⁶ combine with a carbon atom attached thereto to form 1,2,3,4-tetrahydronaphthalene or indan.

[0036] In formula (Ia), X preferably represents CH.

[0037] Preferably, R¹⁸ and R¹⁹, which may be the same or different, represent C₁₋₆ alkoxy optionally substituted by a saturated or unsaturated three- to seven-membered carbocyclic or heterocyclic group.

[0038] Preferably, R¹⁸ and R¹⁹, which may be the same or different, represent C₁₋₆ alkoxy, more preferably methoxy or R³¹-(CH₂)_p-O-, optionally substituted by R¹⁷-(S)m-.

[0039] Preferably, at least one of R²⁰, R²¹, R²² and R²³ represents a halogen atom, preferably a chlorine or fluorine atom, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio, trifluoromethyl, nitro, or amino.


[0040] Preferably, R²⁰ and R²¹, which may be the same or different, represent a halogen atom, more preferably a chlorine or fluorine atom, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio, trifluoromethyl, nitro, or amino, and R²² and R²³ represent a hydrogen atom.

[0041] Preferably, R²¹ and R²², which may be the same or different, represent a halogen atom, more preferably a chlorine or fluorine atom, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio, trifluoromethyl, nitro, or amino, and R²⁰ and R²³ represent a hydrogen atom.

[0042] Preferably, all of R²⁰, R²¹, R²², and R²³ represent a hydrogen atom.

[0043] Preferably, R²⁶ represents thiazolyl or 4-fluorophenyl.

[0044] A group of preferred compounds represented by formula (I) include compounds represented by formula (Ib)





wherein

R^{31} represents a hydrogen atom, a fluorine atom at 2-position, a fluorine atom at 3-position, methoxy at 2-position, methoxy at 3-position, or methyl at 2- and 5-positions,

R^{32} represents methyl, and


R^{33} represents a hydrogen atom, methyl at 1-position, methyl at 2-position, or methyl at 1- and 2-positions.

[0045] Formula (Ib) preferably represents formula (Ib-1) and formula (Ib-2):



wherein R^{31} , R^{32} , and R^{33} are as defined in formula (Ib).

[0046] A group of preferred compounds represented by formula (I) include compounds represented by formula (Ic)




wherein

R^{41} represents a hydrogen atom, a fluorine atom at 2-position, a fluorine atom at 3-position, a chlorine atom at 2-position, a chlorine atom at 3-position, methyl at 2- and 3-positions, methyl at 2- and 5-positions, methoxy at 2-position, methoxy at 3-position, methyl at 2-position, or trifluoromethyl at 2-position,

R^{42} represents methyl,

R^{43} represents a fluorine atom at 4-position, a bromine atom at 3-position, a bromine atom at 4-position, methoxy at 2-position, methoxy at 3-position, methoxy at 4-position, a chlorine atom at 4-position, methyl at 4-position, or nitro at 4-position.

[0047] A group of preferred compounds represented by formula (I) include compounds represented by formula (Id)



wherein

X represents CH or N,

R^{51} represents a hydrogen atom, a fluorine atom at 2-position, a fluorine atom at 3-position, methoxy at 2-position, methoxy at 3-position, or methyl at 2- and 5-positions,


R^{52} represents methyl,

R^{53} represents imidazolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, or isothiazolyl in which one or two hydrogen atoms on the groups are optionally substituted by a halogen atom or C_{1-4} alkyl, and


R^{54} and R^{55} , which may be the same or different, represent a hydrogen atom or C_{1-6} alkyl in which the alkyl group is optionally substituted by hydroxyl; a halogen atom; $-OR^{56}$ wherein R^{56} represents C_{1-4} alkyl; $-NR^{57}R^{58}$ wherein R^{57} and R^{58} , which may be the same or different, represent a hydrogen atom or C_{1-4} alkyl in which the alkyl group is optionally substituted by hydroxyl or $-OR^{59}$ wherein R^{59} represents C_{1-4} alkyl; or a saturated or unsaturated three- to seven-membered carbocyclic or heterocyclic group in which the carbocyclic and heterocyclic groups are optionally substituted by one or two halogen atoms or C_{1-4} alkyl.

[0048] In formula (Id), preferably,

X represents CH, and R^{52} represents




[0049] In formula (Id), more preferably,
X represents CH, R⁵² represents




and
R⁵⁴ and R⁵⁵ represent methyl.

[0050] In formula (Id), more preferably,
X represents CH, R⁵² represents




R⁵⁴ represents methyl, and
R⁵⁵ represents C₁₋₄ alkyl substituted by a saturated or unsaturated five- or six-membered carbocyclic or hetero-
cyclic group.

[0051] In formula (Id), preferably,
X represents CH, and R⁵² represents




[0052] In formula (Id), more preferably,
X represents CH, R⁵² represents




and
R⁵⁴ and R⁵⁵ represent methyl.

[0053] In formula (Id), more preferably,
X represents CH, R⁵² represents




R⁵⁴ represents methyl, and
R⁵⁵ represents C₁₋₄ alkyl substituted by a saturated or unsaturated five- or six-membered carbocyclic or hetero-
cyclic group.

[0054] In formula (Id), preferably,
X represents N, R⁵² represents




[0055] In formula (Id), more preferably,
X represents N, R⁵² represents




and

R⁵⁴ and R⁵⁵ represent methyl.
[0056] In formula (Id), more preferably,
X represents N, R⁵² represents




R⁵⁴ represents methyl, and
R⁵⁵ represents C₁₋₄ alkyl substituted by a saturated or unsaturated five- or six-membered carbocyclic or hetero-
cyclic group.

[0057] In formula (Id), preferably,
X represents N, and R⁵² represents




[0058] In formula (Id), more preferably,
X represents N, R⁵² represents




and

R⁵⁴ and R⁵⁵ represent methyl.
[0059] In formula (Id), more preferably,
X represents N, R⁵² represents



R⁵⁴ represents methyl, and
R⁵⁵ represents C₁₋₄ alkyl substituted by a saturated or unsaturated five- or six-membered carbocyclic or hetero-
cyclic group.

[0060] A group of preferred compounds represented by formula (I) include compounds represented by formula (Ie)



wherein

R⁶⁰¹ represents a hydrogen atom, a fluorine atom at 2-position, a fluorine atom at 3-position, a chlorine atom at
2-position, a chlorine atom at 3-position, methyl at 2- and 3-positions, methyl at 2- and 5-positions, methoxy at 2-po-
sition, methoxy at 3-position, methyl at 2-position, or trifluoromethyl at 2-position,

R⁶⁰² represents methyl,

X represents N or CH₂,

R⁶⁰⁴ and R⁶⁰⁵, which may be the same or different, represent a hydrogen atom or C₁₋₆ alkyl in which the alkyl group is optionally substituted by hydroxyl; a halogen atom; -OR⁶⁰⁶ wherein R⁶⁰⁶ represents C₁₋₄ alkyl; -NR⁶⁰⁷R⁶⁰⁸ wherein R⁶⁰⁷ and R⁶⁰⁸, which may be the same or different, represent a hydrogen atom or C₁₋₄ alkyl in which the alkyl group is optionally substituted by hydroxyl or -OR⁶⁰⁹ wherein R⁶⁰⁹ represents C₁₋₄ alkyl; or a saturated or unsaturated three- to seven-membered carbocyclic or heterocyclic group in which the carbocyclic and heterocyclic groups are optionally substituted by one or two halogen atoms or C₁₋₄ alkyl, and

R⁶¹¹, R⁶¹², R⁶¹³, R⁶¹⁴, and R⁶¹⁵, which may be the same or different, represent a hydrogen atom; hydroxyl; C₁₋₆ alkyl; -OR⁶¹⁶ wherein R⁶¹⁶ represents C₁₋₄ alkyl; a halogen atom; nitro; or -NR⁶¹⁷R⁶¹⁸ wherein R⁶¹⁷ and R⁶¹⁸, which may be the same or different, represent a hydrogen atom or C₁₋₄ alkyl in which the alkyl group is optionally substituted by hydroxyl, -OR⁶¹⁹ wherein R⁶¹⁹ represents C₁₋₄ alkyl, or -NR⁶²⁰R⁶²¹ wherein R⁶²⁰ and R⁶²¹, which may be the same or different, represent a hydrogen atom or C₁₋₄ alkyl.

[0061] Compounds in a preferred embodiment represented by formula (Ie) include

(1) compounds in which X represents CH and all of R⁶¹¹, R⁶¹², R⁶¹³, R⁶¹⁴, and R⁶¹⁵ represent a hydrogen atom, or any one of R⁶¹¹, R⁶¹², R⁶¹³, R⁶¹⁴, and R⁶¹⁵ represents a group other than a hydrogen atom, preferably, C₁₋₆ alkyl, -OR⁶¹⁶, a halogen atom, or nitro, and the remaining groups represent a hydrogen atom,

(2) compounds in which X represents CH and all of R⁶¹¹, R⁶¹², R⁶¹³, R⁶¹⁴, and R⁶¹⁵ represent a hydrogen atom, or any one of R⁶¹¹, R⁶¹², R⁶¹³, R⁶¹⁴, and R⁶¹⁵ represents a group other than a hydrogen atom, preferably, C₁₋₆ alkyl, -OR⁶¹⁶, a halogen atom, or nitro, and the remaining groups represent a hydrogen atom, R⁶⁰⁴ and R⁶⁰⁵ represent methyl,

(3) compounds in which X represents CH and all of R⁶¹¹, R⁶¹², R⁶¹³, R⁶¹⁴, and R⁶¹⁵ represent a hydrogen atom, or any one of R⁶¹¹, R⁶¹², R⁶¹³, R⁶¹⁴, and R⁶¹⁵ represents a group other than a hydrogen atom, preferably, C₁₋₆ alkyl, -OR⁶¹⁶, a halogen atom, or nitro, and the remaining groups represent a hydrogen atom, R⁶⁰⁴ represents methyl and R⁶⁰⁵ represents C₁₋₄ alkyl substituted by a saturated or unsaturated five- or six-membered carbocyclic or heterocyclic group,

(4) compounds in which X represents N and all of R⁶¹¹, R⁶¹², R⁶¹³, R⁶¹⁴, and R⁶¹⁵ represent a hydrogen atom, or any one of R⁶¹¹, R⁶¹², R⁶¹³, R⁶¹⁴, and R⁶¹⁵ represents a group other than a hydrogen atom, preferably, C₁₋₆ alkyl, -OR⁶¹⁶, a halogen atom, or nitro, and the remaining groups represent a hydrogen atom,


(5) compounds in which X represents N and all of R⁶¹¹, R⁶¹², R⁶¹³, R⁶¹⁴, and R⁶¹⁵ represent a hydrogen atom, or any one of R⁶¹¹, R⁶¹², R⁶¹³, R⁶¹⁴, and R⁶¹⁵ represents a group other than a hydrogen atom, preferably, C₁₋₆ alkyl, -OR⁶¹⁶, a halogen atom, or nitro, and the remaining groups represent a hydrogen atom, R⁶⁰⁴ and R⁶⁰⁵ represent methyl, and

(6) compounds in which X represents N and any one of R⁶¹¹, R⁶¹², R⁶¹³, R⁶¹⁴, and R⁶¹⁵ represents a group other than a hydrogen atom, preferably, C₁₋₆ alkyl, -OR⁶¹⁶, a halogen atom, or nitro, and the remaining groups represent a hydrogen atom, R⁶⁰⁴ represents methyl and R⁶⁰⁵ represents C₁₋₄ alkyl substituted by a saturated or unsaturated five- or six-membered carbocyclic or heterocyclic group.

[0062] In each of preferred embodiments (1) to (6) represented by formula (Ie), more preferred embodiments of the compounds in which any one of R⁶¹¹, R⁶¹², R⁶¹³, R⁶¹⁴, and R⁶¹⁵ represents a group other than a hydrogen atom include

- compounds in which R⁶¹¹ represents methoxy and R⁶¹², R⁶¹³, R⁶¹⁴, and R⁶¹⁵ represent a hydrogen atom,
- compounds in which R⁶¹² represents a bromine atom or methoxy and R⁶¹¹, R⁶¹³, R⁶¹⁴, and R⁶¹⁵ represent a hydrogen atom, and
- compounds in which R⁶¹³ represents a bromine atom, a chlorine atom, a fluorine atom, methyl, methoxy, or nitro with R⁶¹¹, R⁶¹², R⁶¹⁴, and R⁶¹⁵ representing a hydrogen atom.

[0063] A group of preferred compounds represented by formula (I) include compounds represented by formula (If)



wherein

X represents CH or N,

R⁷⁰¹ represents a hydrogen atom, a fluorine atom at 2-position, a fluorine atom at 3-position, methoxy at 2-position, methoxy at 3-position, or methyl at 2- and 5-positions,


R⁷⁰² represents C₁₋₄ alkyl,

R⁷⁰³ represents imidazolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, or isothiazolyl in which one or two hydrogen atoms on the groups are optionally substituted by a halogen atom or C₁₋₄ alkyl, and

R⁷⁰⁴ and R⁷⁰⁵, which may be the same or different, represent a hydrogen atom; hydroxyl; nitro; cyano; a halogen atom; -NR⁷⁰⁶R⁷⁰⁷ wherein R⁷⁰⁶ and R⁷⁰⁷, which may be the same or different, represent a hydrogen atom or C₁₋₄ alkyl in which the alkyl group is optionally substituted by hydroxyl, -OR⁷⁰⁸ wherein R⁷⁰⁸ represents C₁₋₄ alkyl, or -NR⁷⁰⁹R⁷¹⁰ wherein R⁷⁰⁹ and R⁷¹⁰, which may be the same or different, represent a hydrogen atom or C₁₋₄ alkyl; -CONR⁷¹¹R⁷¹² wherein R⁷¹¹ and R⁷¹², which may be the same or different, represent a hydrogen atom or C₁₋₄ alkyl in which the alkyl group is optionally substituted by hydroxyl, -OR⁷¹³ wherein R⁷¹³ represents C₁₋₄ alkyl, or -NR⁷¹⁴R⁷¹⁵ wherein R⁷¹⁴ and R⁷¹⁵, which may be the same or different, represent a hydrogen atom or C₁₋₄ alkyl; -COOR⁷¹⁶ wherein R⁷¹⁶ represents a hydrogen atom or C₁₋₄ alkyl in which the alkyl group is optionally substituted by hydroxyl, -OR⁷¹⁷ wherein R⁷¹⁷ represents C₁₋₄ alkyl, or -NR⁷¹⁸R⁷¹⁹ wherein R⁷¹⁸ and R⁷¹⁹, which may be the same or different, represent a hydrogen atom or C₁₋₄ alkyl; C₁₋₆ alkyl; C₂₋₆ alkenyl; C₂₋₆ alkynyl; or C₁₋₆ alkoxy, in which the alkyl, alkenyl, alkynyl, and alkoxy groups are optionally substituted by hydroxyl, a halogen atom, -OR⁷²⁰ in which R⁷²⁰ represents C₁₋₄ alkyl, -NR⁷²¹R⁷²² wherein R⁷²¹ and R⁷²², which may be the same or different, represent a hydrogen atom or C₁₋₄ alkyl in which the alkyl group is optionally substituted by hydroxyl or -OR⁷²³ wherein R⁷²³ represents C₁₋₄ alkyl, or a saturated or unsaturated three- to seven-membered carbocyclic or heterocyclic group in which the carbocyclic and heterocyclic groups are optionally substituted by one or two halogen atoms or C₁₋₄ alkyl.

[0064] In formula (If), preferably,


X represents CH, and R⁷⁰² represents



and, more preferably, in this configuration, R⁷⁰² represents methyl.


[0065] In formula (If), more preferably,

X represents CH, and R⁷⁰² represents



and, more preferably, in this configuration, R⁷⁰² represents methyl, and R⁷⁰⁴ and R⁷⁰⁵ represent methoxy.

[0066] In formula (If), more preferably, X represents CH, and R⁷⁰² represents



and, more preferably, in this configuration, R⁷⁰² represents methyl,

R⁷⁰⁴ represents methoxy, and


R⁷⁰⁵ represents C₁₋₄ alkoxy substituted by a saturated or unsaturated five- or six-membered carbocyclic or heterocyclic group.

[0067] In formula (If), preferably, X represents CH, and R⁷⁰² represents




and, more preferably, in this configuration, R⁷⁰² represents methyl.

[0068] In formula (If), more preferably, X represents CH, and R⁷⁰² represents



and, more preferably, in this configuration, R⁷⁰² represents methyl, and R⁷⁰⁴ and R⁷⁰⁵ represent methoxy.

[0069] In formula (If), more preferably, X represents CH, R⁷⁰² represents




and, more preferably, in this configuration, R⁷⁰² represents methyl, and

R⁷⁰⁴ represents methoxy, and R⁷⁰⁵ represents C₁₋₄ alkoxy substituted by a saturated or unsaturated five- or six-membered carbocyclic or heterocyclic group.

[0070] In formula (If), preferably, X represents N, and R⁷⁰² represents

5




and, more preferably, in this configuration, R⁷⁰² represents methyl.

10

[0071] In formula (If), more preferably,
X represents N, and R⁷⁰² represents

15



20

and, more preferably, in this configuration, R⁷⁰² represents methyl, and
R⁷⁰⁴ and R⁷⁰⁵ represent methoxy.

[0072] In formula (If), more preferably,
X represents N, and R⁷⁰² represents

25



30

and, more preferably, in this configuration, R⁷⁰² represents methyl,


35

R⁷⁰⁴ represents methoxy, and

R⁷⁰⁵ represents C₁₋₄ alkoxy substituted by a saturated or unsaturated five- or six-membered carbocyclic or heterocyclic group.

[0073] In formula (If), preferably,
X represents N, and R⁷⁰² represents

40



45

and, more preferably, in this configuration, R⁷⁰² represents methyl.

[0074] In formula (If), more preferably,
X represents N, and R⁷⁰² represents

50



55

and, more preferably, in this configuration, R⁷⁰² represents methyl, and
R⁷⁰⁴ and R⁷⁰⁵ represent methoxy.


[0075] In formula (If), more preferably,
X represents N, R⁷⁰² represents



and, more preferably, in this configuration, R⁷⁰² represents methyl, and

R⁷⁰⁴ represents methoxy, and R⁷⁰⁵ represents C₁₋₄ alkoxy substituted by a saturated or unsaturated five- or six-membered carbocyclic or heterocyclic group.

[0076] A group of preferred compounds represented by formula (I) include compounds represented by formula (Ig)



wherein

X represents CH or N,

R⁸⁰¹ represents a hydrogen atom, a fluorine atom at 2-position, a fluorine atom at 3-position, a chlorine atom at 2-position, a chlorine atom at 3-position, methyl at 2- and 3-positions, methyl at 2- and 5-positions, methoxy at 2-position, methoxy at 3-position, methyl at 2-position, or trifluoromethyl at 2-position,

R⁸⁰² represents C₁₋₄ alkyl,

R⁸⁰⁴ and R⁸⁰⁵, which may be the same or different, represent a hydrogen atom; hydroxyl; nitro; cyano; a halogen atom; -NR⁸⁰⁶R⁸⁰⁷ wherein R⁸⁰⁶ and R⁸⁰⁷, which may be the same or different, represent a hydrogen atom or C₁₋₄ alkyl in which the alkyl group is optionally substituted by hydroxyl, -OR⁸⁰⁸ wherein R⁸⁰⁸ represents C₁₋₄ alkyl, or -NR⁸⁰⁹R⁸¹⁰ wherein R⁸⁰⁹ and R⁸¹⁰, which may be the same or different, represent a hydrogen atom or C₁₋₄ alkyl; -CONR⁸¹¹R⁸¹² wherein R⁸¹¹ and R⁸¹², which may be the same or different, represent a hydrogen atom or C₁₋₄ alkyl in which the alkyl group is optionally substituted by hydroxyl, -OR⁸¹³ wherein R⁸¹³ represents C₁₋₄ alkyl, or -NR⁸¹⁴R⁸¹⁵ wherein R⁸¹⁴ and R⁸¹⁵, which may be the same or different, represent a hydrogen atom or C₁₋₄ alkyl; -COOR⁸¹⁶ wherein R⁸¹⁶ represents a hydrogen atom or C₁₋₄ alkyl in which the alkyl group is optionally substituted by hydroxyl, -OR⁸¹⁷ wherein R⁸¹⁷ represents C₁₋₄ alkyl, or -NR⁸¹⁸R⁸¹⁹ wherein R⁸¹⁸ and R⁸¹⁹, which may be the same or different, represent a hydrogen atom or C₁₋₄ alkyl; C₁₋₆ alkyl; C₂₋₆ alkenyl; C₂₋₆ alkynyl; or C₁₋₆ alkoxy, in which the alkyl, alkenyl, alkynyl, and alkoxy groups are optionally substituted by hydroxyl, a halogen atom, -OR⁸²⁰ in which R⁸²⁰ represents C₁₋₄ alkyl, -NR⁸²¹R⁸²² wherein R⁸²¹ and R⁸²², which may be the same or different, represent a hydrogen atom or C₁₋₄ alkyl in which the alkyl group is optionally substituted by hydroxyl or -OR⁸²³ wherein R⁸²³ represents C₁₋₄ alkyl, or a saturated or unsaturated three- to seven-membered carbocyclic or heterocyclic group in which the carbocyclic and heterocyclic groups are optionally substituted by one or two halogen atoms or C₁₋₄ alkyl, and

R⁸³¹, R⁸³², R⁸³³, R⁸³⁴, and R⁸³⁵, which may be the same or different, represent a hydrogen atom; hydroxyl; C₁₋₆ alkyl; -OR⁸³⁶ wherein R⁸³⁶ represents C₁₋₄ alkyl; a halogen atom; nitro; or -NR⁸³⁷R⁸³⁸ wherein R⁸³⁷ and R⁸³⁸, which may be the same or different, represent a hydrogen atom or C₁₋₄ alkyl in which the alkyl group is optionally substituted

by hydroxyl, $-OR^{839}$ wherein R^{839} represents C_{1-4} alkyl, or $-NR^{840}R^{841}$ wherein R^{840} and R^{841} , which may be the same or different, represent a hydrogen atom or C_{1-4} alkyl.

[0077] Compounds in a preferred embodiment of formula (Ig) include

- (1) compounds in which X represents CH and all of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represent a hydrogen atom, or any one of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represents a group other than a hydrogen atom, preferably, C_{1-6} alkyl, $-OR^{836}$, a halogen atom, or nitro, and the remaining groups represent a hydrogen atom,
- (2) compounds in which X represents CH and all of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represent a hydrogen atom, or any one of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represents a group other than a hydrogen atom, preferably, C_{1-6} alkyl, $-OR^{836}$, a halogen atom, or nitro, and the remaining groups represent a hydrogen atom, and R^{804} and R^{805} represent methoxy,
- (3) compounds in which X represents CH and all of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represent a hydrogen atom, or any one of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represents a group other than a hydrogen atom, preferably C_{1-6} alkyl, $-OR^{836}$, a halogen atom, or nitro, and the remaining groups represent a hydrogen atom, R^{804} represents methoxy and R^{805} represents C_{1-4} alkoxy substituted by a saturated or unsaturated five- or six-membered carbocyclic or heterocyclic group,
- (4) compounds in which X represents CH, R^{802} represents methyl, and all of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represent a hydrogen atom, or any one of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represents a group other than a hydrogen atom, preferably C_{1-6} alkyl, $-OR^{836}$, a halogen atom, or nitro, and the remaining groups represent a hydrogen atom,
- (5) compounds in which X represents CH, and all of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represent a hydrogen atom, or any one of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represents a group other than a hydrogen atom, preferably C_{1-6} alkyl, $-OR^{836}$, a halogen atom, or nitro, and the remaining groups represent a hydrogen atom, and R^{804} and R^{805} represent methoxy,
- (6) compounds in which X represents CH, and all of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represent a hydrogen atom, or any one of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represents a group other than a hydrogen atom, preferably C_{1-6} alkyl, $-OR^{836}$, a halogen atom, or nitro, and the remaining groups represent a hydrogen atom, R^{804} represents methoxy and R^{805} represents C_{1-4} alkoxy substituted by a saturated or unsaturated five- or six-membered carbocyclic or heterocyclic group,
- (7) compounds in which X represents N, all of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represent a hydrogen atom, or any one of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represents a group other than a hydrogen atom, preferably C_{1-6} alkyl, $-OR^{836}$, a halogen atom, or nitro, and the remaining groups represent a hydrogen atom,
- (8) compounds in which X represents N, all of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represent a hydrogen atom, or any one of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represents a group other than a hydrogen atom, preferably C_{1-6} alkyl, $-OR^{836}$, a halogen atom, or nitro, and the remaining groups represent a hydrogen atom, and R^{804} and R^{805} represent methoxy,
- (9) compounds in which X represent N, all of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represent a hydrogen atom, or any one of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represents a group other than a hydrogen atom, preferably C_{1-6} alkyl, $-OR^{836}$, a halogen atom, or nitro, and the remaining groups represent a hydrogen atom, R^{804} represents methoxy, and R^{805} represents C_{1-4} alkoxy substituted by a saturated or unsaturated five- or six-membered carbocyclic or heterocyclic group,
- (10) compounds in which X represents N, R^{802} represents methyl, all of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represent a hydrogen atom, or any one of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represents a group other than a hydrogen atom, preferably C_{1-6} alkyl, $-OR^{836}$, a halogen atom, or nitro, and the remaining groups represent a hydrogen atom,
- (11) compounds in which X represents N, all of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represent a hydrogen atom, or any one of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represents a group other than a hydrogen atom, preferably C_{1-6} alkyl, $-OR^{836}$, a halogen atom, or nitro, and the remaining groups represent a hydrogen atom, and R^{804} and R^{805} represent methoxy, and
- (12) compounds in which X represents N, all of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represent a hydrogen atom, or any one of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represents a group other than a hydrogen atom, preferably C_{1-6} alkyl, $-OR^{836}$, a halogen atom, or nitro, and the remaining groups represent a hydrogen atom, R^{804} represents methoxy, and R^{805} represents C_{1-4} alkoxy substituted by a saturated or unsaturated five- or six-membered carbocyclic or heterocyclic group.

[0078] In each of preferred embodiments (1) to (12) of formula (Ig), more preferred embodiments of the compounds in which any one of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represents a group other than a hydrogen atom include

- compounds in which R^{831} represents methoxy and R^{832} , R^{833} , R^{834} , and R^{835} represent a hydrogen atom,
- compounds in which R^{832} represents a bromine atom or methoxy and R^{831} , R^{833} , R^{834} , and R^{835} represent a

hydrogen atom, and

- compounds in which R^{833} represents a bromine atom, a chlorine atom, a fluorine atom, methyl, methoxy, or nitro, with R^{831} , R^{832} , R^{834} , and R^{835} representing a hydrogen atom.

[0079] Examples of preferred compounds of the present invention include compounds described in Examples 1 to 38 and 41 to 105.

[0080] Examples of more preferred compounds according to the present invention include the following compounds:

- (16) N-{4-[(6,7-dimethoxy-4-quinolyl)oxy]-2-methoxyphenyl}-N'-[1-(4-fluorophenyl)ethyl]urea;
- (17) N-{4-[(6,7-dimethoxy-4-quinolyl)oxy]-2-methoxyphenyl}-N'-[(1S)-1-(4-fluorophenyl)ethyl]urea;
- (18) N-{4-[(6,7-dimethoxy-4-quinolyl)oxy]-2-methoxyphenyl}-N'-[(1R)-1-(4-fluorophenyl)ethyl]urea;
- (74) N-{4-[(6,7-dimethoxy-4-quinolyl)oxy]-2-methoxyphenyl}-N'-[1-(1,3-thiazol-2-yl)ethyl]urea;
- (75) N-{4-[(6,7-dimethoxy-4-quinolyl)oxy]-2-methoxyphenyl}-N'-[(1S)-1-(1,3-thiazol-2-yl)ethyl]urea; and
- (76) N-{4-[(6,7-dimethoxy-4-quinolyl)oxy]-2-methoxyphenyl}-N'-[(1R)-1-(1,3-thiazol-2-yl)ethyl]urea.

[0081] Compounds according to the present invention may form pharmaceutically acceptable salts thereof. Preferred examples of such salts include: alkali metal or alkaline earth metal salts such as sodium salts, potassium salts or calcium salts; hydrohalogenic acid salts such as hydrofluoride salts, hydrochloride salts, hydrobromide salts, or hydroiodide salts; inorganic acid salts such as nitric acid salts, perchloric acid salts, sulfuric acid salts, or phosphoric acid salts; lower alkylsulfonic acid salts such as methanesulfonic acid salts, trifluoromethanesulfonic acid salts, or ethanesulfonic acid salts; arylsulfonic acid salts such as benzenesulfonic acid salts or p-toluenesulfonic acid salts; organic acid salts such as fumaric acid salts, succinic acid salts, citric acid salts, tartaric acid salts, oxalic acid salts, maleic acid salts, acetic acid salts, malic acid salts, lactic acid salts, or ascorbic acid salts; and amino acid salts such as glycine salts, phenylalanine salts, glutamic acid salts, or aspartic acid salts.


[0082] Pharmaceutically acceptable solvates of the compounds according to the present invention include, for example, hydrates, alcoholates, for example, ethanolates, and etherates.

[0083] One or more enantiomeric carbon atoms, which form enantiomer configuration, may exist in the compounds according to the present invention. The compounds according to the present invention include all enantiomers.

Production of compounds

[0084] Compounds according to the present invention may be produced, for example, according to scheme 1, scheme 2 scheme 3, and scheme 4.

Scheme 1



[0085] In the formula, R represents C_{1-6} alkyl or the like; and $R^1, R^2, R^3, R^4, R^5, R^6, R^7, R^8$, and X are as defined in formula (I).

[0086] Starting compounds necessary for the synthesis of the compounds according to the present invention are commercially available or can be easily produced by a conventional method. For example, 4-chloroquinoline derivatives may be synthesized by a conventional method described, for example, in Org. Synth. Col. Vol. 3, 272 (1955), Acta Chim. Hung., 112, 241 (1983) or WO 98/47873.


[0087] Alternatively, 4-chloroquinazolinone derivatives may be produced by first (1) reacting a benzoic ester with formamide to give a quinazolinone derivative and subsequently (2) heating the 4-quinazolinone derivatives using toluene or sulfolane as a solvent in the presence of phosphorus oxychloride. The quinazolinone derivative may be synthesized by reacting a benzoic ester, sodium methoxide, and formamide in the presence of a solvent such as N,N-dimethylformamide or methanol.

[0088] Next, a 4-(aminophenoxy)quinoline derivative or a corresponding quinazolinone derivative is produced by reacting nitrophenol with the 4-chloroquinoline derivative or corresponding quinazolinone derivative in the presence or absence of a suitable solvent to synthesize a 4-(nitrophenoxy)quinoline derivative or a corresponding quinazolinone derivative.

rivative, then conducting stirring in a suitable solvent, for example, N,N-dimethyl formamide, in the presence of a catalyst, for example, palladium hydroxide-carbon or palladium-carbon, under a hydrogen atmosphere. Alternatively, the 4-(aminophenoxy)quinoline derivative or a corresponding quinazoline derivative may also be produced by reacting amino phenol with a 4-chloroquinoline derivative or a corresponding quinazoline derivative in the presence of a base, for example, sodium hydride.

[0089] Alternatively, the 4-(aminophenoxy)quinazoline derivative may be produced by dissolving amino phenol in an aqueous sodium hydroxide solution and subjecting the solution to a two-phase reaction with a solution of the 4-chloroquinazoline derivative in an organic solvent in the presence of a phase transfer catalyst, for example, tetra-n-butylammonium bromide, or in the absence of the catalyst.


Scheme 2



wherein R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, R¹¹, R¹², R¹³, and X are as defined in formula (I).

[0090] A urea derivative in which R⁹ and/or R¹⁰ represent a hydrogen atom may be produced according to a conventional method, i.e., by adding triphosgene or a chloroformic ester to a 4-(aminophenoxy)quinoline derivative prepared in scheme 1 or a corresponding quinazoline derivative and then reacting the mixture with a suitable amine derivative (R¹¹R¹²R¹³CNH₂) in the presence of a base, for example, triethylamine (scheme 2).

Scheme 3




wherein R' represents hydroxyl, C₁₋₄ alkyl or the like; and R¹², R¹³, and X are as defined in formula (I).

[0091] The amine derivative (R¹¹R¹²R¹³CNH₂) may be produced, for example, by reacting a commercially available acylated aryl compound or a heteroaryl compound with a suitable primary amine, for example, hydroxylamine, for conversion to an imine derivative and then treating the imine derivative with a suitable reducing agent, for example, zinc (scheme 3).

[0092] R⁹ and R¹⁰ can be introduced by reacting a urea derivative, in which R⁹ and/or R¹⁰ represent a hydrogen atom, with a suitable alkylating agent (R⁹Hal or R¹⁰Hal wherein Hal represents a halogen atom) in the presence of a base, for example, sodium hydride.

Scheme 4



wherein PG represents a protective group of hydroxyl; R^3 represents C_{1-6} alkyl, wherein the alkyl group is optionally substituted by the same substituent as the substituent of alkoxy represented by R^3 ; and R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , R^{11} , R^{12} , and R^{13} are as defined in formula (I).

[0093] The derivative having a specific substituent at the 7-position of the quinoline ring may be produced, for example, by scheme 4.

[0094] A nitro group may be introduced by first reacting a commercially available 4'-hydroxyacetophenone derivative with a suitable substituent, for example, benzyl, to protect hydroxyl, and then reacting the reaction product with a nitrating agent, for example, nitric acid-acetic acid.

[0095] Next, a 4-chloroquinoline derivative may be produced by reducing the nitro group to give an amino group, then reacting the amino group with a formic ester in the presence of a base to give a quinolone which was then reacted with a chlorinating agent, for example, phosphorus oxychloride.

[0096] An 4-(aminophenoxy)quinoline derivative is produced by reacting an aminophenol with the 4-chloroquinoline derivative prepared above in the presence of a base, for example, sodium hydride. Next, a urea derivative may be produced by a conventional method, i.e., by adding triphosgene or a chloroformic ester in the presence of a base, for example, triethylamine, and then reacting the mixture with a suitable amine derivative ($R^{11}R^{12}R^{13}CNH_2$).

[0097] The compound of the present invention having alkoxy at the 7-position of the quinoline ring can be produced by removing the protective group (PG) of hydroxyl at the 7-position of the quinoline ring and reacting the deprotected compound with an alkyl halide in the presence of a base or by treating an alcohol derivative according to a conventional method, for example, Mitsunobu reaction.

[0098] The alkyl halide used in the substitution reaction is commercially available or alternatively may be produced according to a method described, for example, in J. Am. Chem. Soc., 1945, 67, 736.

[0099] The alcohol derivative used in the substitution reaction is commercially available, or alternatively may be produced by a method, for example, described in J. Antibiot. (1993), 46 (1), 177 and Ann. Pharm. Fr. 1997, 35, 503.

[0100] The derivative having a specific substituent at the 6-position of the quinoline ring may be produced in the

same manner as in scheme 4, except that a 3'-hydroxyacetophenone derivative is used as a starting compound.

[0101] The derivative having a specific substituent at the 7-position of the quinazoline ring may be produced in the same manner as in scheme 4, except that a 4-hydroxybenzoic ester derivative is used as a starting compound and formamide instead of the formic ester.

[0102] The derivative having a specific substituent at the 6-position of the quinazoline ring may be produced in the same manner as in scheme 4, except that a 3-hydroxybenzoic ester derivative is used as a starting compound and formamide instead of the formic ester.

[0103] The compound represented by formula (I) wherein Z represents S may be produced according to scheme 1 by reacting an aminothiophenol derivative with a 4-chloroquinoline derivative or a corresponding quinazoline derivative in a suitable solvent, for example, chlorobenzene, to give a 4-(quinolylsulfanyl)aniline derivative or a 4-(quinazoliny-sulfanyl)aniline derivative which is then reacted with an amine derivative according to scheme 2.

Use of compounds/pharmaceutical composition

[0104] Bone is a tissue which is maintained by repeating bone resorption and bone formation, and cells responsible for a bone resorption function are osteoclasts. Various factors are known as factors for accelerating differentiation of osteoclasts in vivo. RANKL which has been recently identified as a differentiation-inducing factor of osteoclasts may be mentioned as a representative example of such factors. RANKL induces differentiation of osteoclasts from splenic cells and marrow cells in vitro. In this case, the presence of a macrophage colony stimulating factor (hereinafter referred to as "M-CSF") is indispensable. On the other hand, that M-CSF is indispensable in vivo is apparent from the fact that, upon deletion of osteoclasts, op/op mice, i.e., animals which do not produce normal M-CSF, exhibit osteopetrosis (Yoshida H, et al., Nature, 345, 442-444 (1990) "The Murine Mutation Osteopetrosis is in the Coding Region of the Macrophage Colony Stimulating Factor Gene"), and the fact that the administration of activated M-CSF to the mice can cure osteopetrosis. Osteoclasts are important cells responsible for bone resorption in maintaining bone tissue, but on the other hand, in diseases such as bone metastasis of malignant tumors including breast cancer, prostatic cancer, and lung cancer, multiple myeloma, osteoporosis, Behcet's disease, rheumatoid arthritis or other diseases, enhanced bone resorption of osteoclasts in bone tissue is causative of worsening of condition of diseases. Therefore, these clinical conditions and cancer pain caused by bone metastasis can be ameliorated by inhibiting the differentiation of osteoclasts to reduce the number of osteoclasts. Osteoclasts are derived from monocytoïd cells which are hemocytes. In precursor cells thereof, signals which promote differentiation of osteoclasts are transmitted through an M-CSF receptor (hereinafter referred to as "c-fms"). It is known that c-fms is receptor tyrosine kinase and binding of M-CSF thereto leads to activation of tyrosine kinase on the receptor. The receptor tyrosine kinase functions to transmit signals through autophosphorylation of the receptor per se. Therefore, differentiation induction to osteoclasts can be inhibited by inhibiting the autophosphorylation of c-fms. As a result, the number of osteoclasts in vivo can be reduced.

[0105] Compounds according to the present invention inhibited the autophosphorylation of mouse c-fms which occurs upon stimulation of macrophage-like cell line BAC-1.2F5, capable of stably expressing mouse c-fms in vitro, by M-CSF (see Pharmacological Test Example 1).

[0106] Compounds according to the present invention exhibited concentration-dependent inhibition of M-CSF-dependent cell growth of BAC-1.2F5 in vitro (see Pharmacological Text Example 2).

[0107] Further, compounds according to the present invention exhibited concentration-dependent inhibition of differentiation of osteoclasts in vitro (see Pharmacological Text Example 3).

[0108] Compounds according to the present invention inhibited the expansion of bone resorption area involved in bone metastasis of melanoma in a nude mouse bone metastasis model and in a nude rat bone metastasis model (see Pharmacological Text Examples 4 and 5).

[0109] Furthermore, compounds according to the present invention inhibited differentiation of osteoclasts in a rat osteoporosis model (see Pharmacological Text Example 6).

[0110] Accordingly, the compounds according to the present invention can be used for the therapy and prophylaxis of a disease for which the inhibition of M-CSF receptor autophosphorylation is effective therapeutically.

[0111] Diseases for which the inhibition of M-CSF receptor autophosphorylation is effective therapeutically include bone metastasis of malignant tumors including breast cancer, prostatic cancer, and lung cancer; multiple myeloma; osteoporosis; Behcet's disease; and rheumatoid arthritis.

[0112] The compounds according to the present invention can be administered to human and non-human animals orally or parenterally by administration routes, for example, intravenous administration, intramuscular administration, subcutaneous administration, rectal administration, or percutaneous administration. Therefore, the pharmaceutical composition comprising as an active ingredient the compound according to the present invention is formulated into suitable dosage forms according to the administration routes.

[0113] Specifically, oral preparations include tablets, capsules, powders, granules, and syrups, and parental preparations include injections, suppositories, tapes, and ointments.

[0114] These various preparations may be prepared by conventional methods, for example, with commonly used excipients, disintegrants, binders, lubricants, colorants, and diluents.

[0115] Excipients include, for example, lactose, glucose, corn starch, sorbit, and crystalline cellulose. Disintegrants include, for example, starch, sodium alginate, gelatin powder, calcium carbonate, calcium citrate, and dextrin. Binders include, for example, dimethylcellulose, polyvinyl alcohol, polyvinyl ether, methylcellulose, ethylcellulose, gum arabic, gelatin, hydroxypropylcellulose, and polyvinyl pyrrolidone. Lubricants include, for example, talc, magnesium stearate, polyethylene glycol, and hydrogenated vegetable oils.

[0116] In preparing the injections, if necessary, for example, buffers, pH adjusters, stabilizers, tonicity agents, and preservatives may be added.

[0117] The content of the compound according to the present invention in the pharmaceutical composition according to the present invention may vary depending upon the dosage form. In general, however, the content is 0.5 to 50% by weight, preferably 1 to 20% by weight, based on the whole composition.

[0118] The dose may be appropriately determined in consideration of, for example, the age, weight, sex, difference in diseases, and severity of condition of individual patients, for example, in the range of 0.01 to 100 mg/kg, preferably 0.1 to 50 mg/kg. This dose is administered at a time daily or divided doses of several times daily.

[0119] The compound according to the present invention may be administered in combination with other medicament. In this case, the compound according to the present invention may be administered simultaneously with or after or before the administration of other medicament. For example, when the target disease is bone metastasis of malignant tumors such as breast cancer, prostatic cancer, and lung cancer, or osteolysis caused by malignant tumors such as multiple myeloma, it is considered that the action of the compound according to the present invention on precursor cells of osteoclasts reduces osteoclasts to inhibit osteolysis and subsequent administration of a carcinostatic agent can efficiently inhibit the growth of the malignant tumor within the bone. The kind of the carcinostatic agent, dosage interval and the like can be determined depending upon the kind of cancer and the condition of the patient. Diseases other than malignant tumors can also be similarly treated.

[0120] According to the present invention, the contact of the compound of the present invention with precursor cells of osteoclasts can be conducted, for example, by systemic administration, for example, intravenous administration or oral administration, topical administration, for example, dermal administration or intraarticular administration, drug targeting using a carrier, for example, liposome, lipid microsphere, or polymerized pharmaceutical preparation.

[0121] According to the present invention, there is provided use of the compound according to the present invention, for the manufacture of an agent for use in the treatment and prevention of a disease for which the inhibition of M-CSF receptor autophosphorylation is effective therapeutically.

[0122] Further, according to the present invention, there is provided a method for treating and preventing a disease for which the inhibition of M-CSF receptor autophosphorylation is effective therapeutically, said method comprising the step of administering a therapeutically or prophylactically effective amount of the compound according to the present invention together with a pharmaceutically acceptable carrier to a mammal.

Selective inhibition of c-fms

[0123] The present invention provides a compound which can selectively inhibit c-fms rather than KDR (see Pharmacological Test Example 7). A group of such compounds include compounds of formula (Ib), formula (Ic), formula (Id), formula (Ie), formula (If), and formula (Ig), particularly, compounds of Examples 17, 74, 75, and 76.

[0124] It is known that proteins including tyrosine kinase domains play a very important role, for example, in signal transmission in vivo and there are many kinds of such proteins. In general, "selective" action of the medicament on the target site is considered to reduce side effects. Therefore, the use of the inhibitor capable of selectively acting on M-CSF receptor (c-fms) tyrosine kinase as the target of the compound according to the present invention is considered to reduce the side effect.

[0125] In addition to c-fms, VEGF receptor (KDR) is known as a protein containing tyrosine kinase domain. It has been proven that the inhibition of the phosphorylation of KDR can inhibit angiogenesis, and a KDR phosphorylation inhibitor is regarded as useful for the treatment of a disease for which the inhibition of angiogenesis is desired. On the other hand, angiogenesis is desired in organs such as kidney. For example, in a progressive renal failure model in rats, there is a report that repair of blood capillaries by VEGF exhibits therapeutic effect (Yukinari Masuda, et al: VEGF Accelerates Glomerular Repair in GN. Am J Pathol, 159: 599-608, 2001). Therefore, KDR phosphorylation inhibitory activity in organs for which angiogenesis is desired possibly has adverse effect on the function of the organs. In particular, in the case of therapy of bone metastasis of cancers or therapy of osteoporosis in which medication is carried out over an extended time, there is a fear of this adverse effect.

[0126] For the above reason, it can be said that, among c-fms inhibitors, compounds which selectively inhibit c-fms rather than KDR is more advantageous from the viewpoint of lower possibility of sacrificing the function of an organ for which angiogenesis is desired.

EXAMPLES

[0127] The present invention is further illustrated by the following Examples that are not intended as a limitation of the invention.

Production Example 1: 1-(1,3-Thiazol-2-yl)-1-ethanone oxime (mixture of isomers)

[0128] Commercially available 2-acetyl thiazole (1.27 g) and commercially available hydroxylamine hydrochloride (830 mg) were dissolved in ethanol (40 ml) to prepare a solution. Pyridine (0.97 ml) was added to the solution, and the mixture was stirred at room temperature overnight. Water was added to the reaction solution, and the mixture was extracted three times with ethyl acetate. The organic layer was washed with water and was dried over sodium sulfate. The solvent was removed by evaporation and was dried by means of a vacuum pump to give the title compound (1.25 g, 88%).

[0129] $^1\text{H-NMR}$ (CDCl_3): δ 2.43 (s, 3H), 2.49 (s, 3H), 7.31 (d, $J = 3.2$ Hz, 1H), 7.60 (d, $J = 3.4$ Hz, 1H), 7.85 (d, $J = 3.2$ Hz, 1H), 8.01 (d, $J = 3.2$ Hz, 1H)

Production Example 2: 1-(1,3-Thiazol-2-yl)-1-ethylamine

[0130] 1-(1,3-Thiazol-2-yl)-1-ethanone oxime (710 mg) was dissolved in methanol (10 ml) to prepare a solution. Ammonium chloride (1.35 g) and zinc (powder) (6.5 g) were added to the solution, and the mixture was heated under reflux for 2 hr. Thereafter, 10% sodium hydroxide (10 ml) was added thereto, and the mixture was extracted three times with chloroform. The organic layer was washed with saturated brine and was dried over sodium sulfate. The solvent was removed by evaporation and was dried by means of a vacuum pump to give the title compound (580 mg, 91%).

[0131] $^1\text{H-NMR}$ (CDCl_3): δ 1.55 (d, $J = 6.8$ Hz, 3H), 4.43 (q, $J = 6.8$ Hz, 3H), 7.24 (d, $J = 3.4$ Hz, 1H), 7.71 (d, $J = 3.2$ Hz, 1H)

Production Example 3: 1-(1,3-Thiazol-2-yl)-1-ethylamine (optically active substance)

[0132] 1-(1,3-Thiazol-2-yl)-1-ethylamine (racemate) (2.0 g) was dissolved in dichloromethane (20 ml) to prepare a solution. Triethylamine (2.5 ml) was added to the solution, and the mixture was cooled to 0°C . Di-tert-butyl dicarbonate (3.7 g) was added thereto, and the mixture was stirred at room temperature for 3 hr. Water (40 ml) was added thereto, and the mixture was extracted three times with chloroform. The organic layer was washed with saturated brine and was dried over sodium sulfate. The solvent was removed, and the residue was purified by chromatography on silica gel using hexane/ethyl acetate for development to give tert-butyl N-[1-(1,3-thiazol-2-yl)ethyl]carbamate (racemate) (2.8 g, 62%).

[0133] $^1\text{H-NMR}$ (CDCl_3): δ 1.46 (s, 9H), 1.60 (d, $J = 6.8$ Hz, 3H), 5.12 (br, 1H), 5.25 (br, 1H), 7.25 (dd, $J = 0.5, 3.2$ Hz, 1H), 7.70 (dd, $J = 0.5, 3.2$ Hz, 1H)

[0134] Tert-butyl N-[1-(1,3-thiazol-2-yl)ethyl]carbamate (racemate) prepared above was resolved into optical isomers with an optical isomer resolution column (Daicel Chemical Industries, Ltd.). Optically active N-[1-(1,3-thiazol-2-yl)ethyl]carbamate (400 mg) was dissolved in dichloromethane (10 ml) to prepare a solution which was then cooled to 0°C . Trifluoroacetic acid (4 ml) was added to the cooled solution, and the temperature of the mixture was then raised to room temperature before the mixture was stirred for 90 min. Thereafter, 10% aqueous sodium hydroxide solution was added thereto, and, after it was confirmed that the solution was rendered basic, the solution was stirred at room temperature for 30 min and was then extracted three times with chloroform. The organic layer was washed with saturated brine and was dried over sodium sulfate. The solvent was removed by evaporation and was dried by means of a vacuum pump to quantitatively give the title compound.

Production Example 4: 1-(5-Methyl-1,3-thiazol-2-yl)-1-ethanone

[0135] Commercially available 5-methyl-1,3-thiazole (218 mg) was dissolved in tetrahydrofuran (5 ml) to prepare a solution which was then cooled to -78°C . A hexane solution (1.56 M) (1.4 ml) of n-butyllithium was slowly added to the cooled solution over a period of 10 min, and the mixture was stirred at -78°C for 2 hr. N-Methoxy-N-methylacetamide (206 mg) was dissolved in tetrahydrofuran (2 ml) to prepare a solution which was then slowly added thereto over a period of 10 min, followed by stirring at -78°C for 2 hr. The cooling bath was removed, an aqueous saturated ammonium chloride solution (5 ml) was added, and the mixture was stirred for 30 min. Further, water (5 ml) was added thereto, and the reaction solution was extracted with ethyl acetate. The organic layer was washed with saturated brine and was dried over sodium sulfate. The solvent was removed, and the residue was purified by chromatography on silica gel using hexane/ethyl acetate for development to give the title compound (267 mg, 86%).

[0136] $^1\text{H-NMR}$ (CDCl_3): δ 2.56 (d, $J = 1.0$ Hz, 3H), 2.67 (s, 3H), 7.65 (d, $J = 1.0$ Hz, 1H)

Production Example 5: 1-(4,5-Dimethyl-1,3-thiazol-2-yl)-1-ethanone

[0137] Commercially available 4,5-dimethyl-1,3-thiazole (453 mg) was dissolved in tetrahydrofuran (10 ml) to prepare a solution which was then cooled to -78°C . A hexane solution (1.56 M) (2.8 ml) of *n*-butyllithium was slowly added over a period of 10 min, and the mixture was stirred at -78°C for 2 hr. *N*-Methoxy-*N*-methylacetamide (454 mg) was dissolved in tetrahydrofuran (2 ml) to prepare a solution which was then slowly added over a period of 10 min, and the mixture was stirred at -78°C for 2 hr. The cooling bath was removed, an aqueous saturated ammonium chloride solution (10 ml) was added, and the mixture was stirred for 30 min. Further, water (10 ml) was added, and the reaction solution was extracted with ethyl acetate. The organic layer was washed with saturated brine and was dried over sodium sulfate. The solvent was removed, and the residue was purified by chromatography on silica gel using hexane/ethyl acetate for development to give the title compound (609 mg, 98%).

[0138] $^1\text{H-NMR}$ (CDCl_3): δ 2.40 (d, $J = 0.7$ Hz, 3H), 2.44 (s, 3H), 2.64 (d, $J = 0.7$ Hz, 3H)

Example 1: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]phenyl]-N'-[1-(4-fluorophenyl)ethyl]urea

[0139] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]aniline (20 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(4-fluorophenyl)ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The reaction solution was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (2 mg, yield 8%).

[0140] $^1\text{H-NMR}$ (CDCl_3): 1.49 (3H, d, $J = 6.83$ Hz), 4.04 (3H, s), 4.06 (3H, s), 4.92 - 4.99 (1H, m), 6.42 (1H, d, $J = 5.34$ Hz), 7.02 (2H, dd, $J = 8.54, 8.54$ Hz), 7.07 (2H, d, $J = 8.78$ Hz), 7.33 (2H, dd, $J = 8.54, 5.37$ Hz), 7.37 (1H, s), 7.46 (2H, d, $J = 8.78$ Hz), 7.57 (1H, s), 8.39 (1H, d, $J = 5.34$ Hz)

[0141] Mass spectrometric value (ESI-MS, m/z): 460 ($M-1$)

Example 2: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]phenyl]-N'-[(1S)-1-(4-fluorophenyl)ethyl]urea

[0142] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]aniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1S)-1-(4-fluorophenyl)ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (33 mg, yield 71%).

[0143] $^1\text{H-NMR}$ (CDCl_3): δ 1.49 (3H, d, $J = 6.83$ Hz), 4.04 (3H, s), 4.04 (3H, s), 4.94 - 4.99 (1H, m), 6.42 (1H, d, $J = 5.34$ Hz), 7.04 (2H, dd, $J = 8.78, 8.78$ Hz), 7.09 (2H, d, $J = 8.54$ Hz), 7.33 (2H, dd, $J = 8.54, 5.37$ Hz), 7.37 (1H, s), 7.46 (2H, d, $J = 8.78$ Hz), 7.57 (1H, s), 8.42 (1H, d, $J = 5.10$ Hz)

[0144] Mass spectrometric value (ESI-MS, m/z): 462 ($M^+ + 1$)

[0145] $[\alpha]_D^{24} - 27.5^\circ$ (c0.5, CHCl_3)

Example 3: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]phenyl]-N'-[(1R)-1-(4-fluorophenyl)ethyl]urea

[0146] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]aniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1R)-1-(4-fluorophenyl)ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (28 mg, yield 60%).

[0147] $^1\text{H-NMR}$ (CDCl_3): δ 1.51 (3H, d, $J = 6.83$ Hz), 4.04 (3H, s), 4.06 (3H, s), 4.94 - 4.99 (1H, m), 6.42 (1H, d, $J = 5.34$ Hz), 7.02 (2H, dd, $J = 8.54, 8.54$ Hz), 7.07 (2H, d, $J = 8.78$ Hz), 7.33 (2H, dd, $J = 8.54, 5.37$ Hz), 7.39 (1H, s), 7.47 (2H, d, $J = 8.78$ Hz), 7.57 (1H, s), 8.41 (1H, d, $J = 5.10$ Hz)

[0148] Mass spectrometric value (ESI-MS, m/z): 462 ($M^+ + 1$)

[0149] $[\alpha]_D^{24} + 27.1^\circ$ (c0.5, CHCl_3)

Example 4: N-(2-Chloro-4-[(6,7-dimethoxy-4-quinolyl)oxy]phenyl)-N'-[1-(4-fluorophenyl)ethyl]urea

[0150] 2-Chloro-4-[(6,7-dimethoxy-4-quinolyl)oxy]aniline (82 mg) was dissolved in chloroform (2 ml) and triethylamine (0.2 ml) to prepare a solution. A solution of triphosgene (70 mg) in chloroform (0.4 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(4-fluorophenyl)ethylamine (70 mg) in chloroform (0.4 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (40 mg, yield 32%).

[0151] ¹H-NMR (CDCl₃): 1.54 (3H, d, J = 6.83 Hz), 4.03 (3H, s), 4.04 (3H, s), 4.96 (1H, m), 5.21 (1H, d, J = 6.59 Hz), 6.45 (1H, d, J = 5.37 Hz), 6.77 (1H, s), 7.03 - 7.10 (3H, overlapped), 7.17 (1H, d, J = 2.68 Hz), 7.36 (2H, dd, J = 6.59, 3.42 Hz), 7.42 (1H, s), 7.49 (1H, s), 8.26 (1H, d, J = 9.03 Hz), 8.48 (1H, d, J = 5.37 Hz)

[0152] Mass spectrometric value (ESI-MS, m/z): 494 (M-1)

Example 5: N-(2-Chloro-4-[(6,7-dimethoxy-4-quinolyl)oxy]phenyl)-N'-[(1S)-1-(4-fluorophenyl)ethyl]urea

[0153] 2-Chloro-4-[(6,7-dimethoxy-4-quinolyl)oxy]aniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1S)-1-(4-fluorophenyl)ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (9 mg, yield 21%).

[0154] ¹H-NMR (CDCl₃): 1.55 (3H, d, J = 6.83 Hz), 4.04 (3H, s), 4.04 (3H, s), 4.92 - 4.98 (1H, m), 5.07 (1H, d, J = 6.59 Hz), 6.45 (1H, d, J = 5.12 Hz), 6.71 (1H, s), 7.04 - 7.10 (3H, overlapped), 7.18 (1H, d, J = 2.68 Hz), 7.38 (2H, dd, J = 8.78, 5.34 Hz), 7.42 (1H, s), 7.50 (1H, s), 8.26 (1H, d, J = 9.03 Hz), 8.49 (1H, d, J = 5.37 Hz)

[0155] Mass spectrometric value (ESI-MS, m/z): 496 (M+1)

[0156] [α]_D²⁴-32.5° (c0.25, CHCl₃)

Example 6: N-(2-Chloro-4-[(6,7-dimethoxy-4-quinolyl)oxy]phenyl)-N'-[(1R)-1-(4-fluorophenyl)ethyl]urea

[0157] 2-Chloro-4-[(6,7-dimethoxy-4-quinolyl)oxy]aniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1R)-1-(4-fluorophenyl)ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (17 mg, yield 37%).

[0158] ¹H-NMR (CDCl₃): 1.55 (3H, d, J = 6.83 Hz), 4.04 (3H, s), 4.04 (3H, s), 4.92 - 4.98 (1H, m), 5.03 (1H, d, J = 6.83 Hz), 6.45 (1H, d, J = 5.37 Hz), 6.69 (1H, s), 7.03 - 7.10 (3H, overlapped), 7.18 (1H, d, J = 2.68 Hz), 7.38 (2H, dd, J = 5.37, 8.78 Hz), 7.42 (1H, s), 7.49 (1H, s), 8.26 (1H, d, J = 9.03 Hz), 8.49 (1H, d, J = 5.37 Hz)

[0159] Mass spectrometric value (ESI-MS, m/z): 496 (M+1)

[0160] [α]_D²⁵+27.9° (c0.25, CHCl₃)

Example 7: N-(4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2,5-dimethylphenyl)-N'-[1-(4-fluorophenyl)ethyl]urea

[0161] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2,5-dimethoxyaniline (81 mg) was dissolved in chloroform (2 ml) and triethylamine (0.2 ml) to prepare a solution. A solution of triphosgene (70 mg) in chloroform (0.4 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(4-fluorophenyl)ethylamine (35 mg) in chloroform (0.4 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (41 mg, yield 33%).

[0162] ¹H-NMR (CDCl₃): 1.49 (3H, d, J = 6.83 Hz), 2.09 (3H, s), 2.12 (3H, s), 4.04 (3H, s), 4.05 (3H, s), 4.97 (1H, m), 5.05 (1H, d, J = 6.80 Hz), 6.09 (1H, s), 6.27 (1H, d, J = 5.37 Hz), 6.91 (1H, s), 7.02 (2H, dd, J = 8.52, 6.36 Hz), 7.33 (2H, dd, J = 8.52, 5.12 Hz), 7.42 (1H, s), 7.52 (1H, s), 7.57 (1H, s), 8.43 (1H, d, J = 5.37 Hz)

[0163] Mass spectrometric value (ESI-MS, m/z): 488 (M-1)

Example 8: N-(4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2,5-dimethylphenyl)-N'-[(1S)-1-(4-fluorophenyl)ethyl]urea

[0164] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2,5-dimethoxyaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to

the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1S)-1-(4-fluorophenyl) ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (35 mg, yield 77%).

[0165] $^1\text{H-NMR}$ (CDCl_3): 1.49 (3H, d, $J = 6.34$ Hz), 2.09 (3H, s), 2.13 (3H, s), 4.05 (3H, s), 4.05 (3H, s), 4.95 - 5.00 (2H, m), 6.28 (1H, d, $J = 5.37$ Hz), 6.92 (1H, s), 7.04 (2H, dd, $J = 8.78, 8.78$ Hz), 7.33 (2H, dd, $J = 8.54, 5.12$ Hz), 7.42 (1H, s), 7.51 (1H, s), 7.57 (1H, s), 8.44 (1H, d, $J = 5.12$ Hz)

[0166] Mass spectrometric value (ESI-MS, m/z): 490 ($M+1$)

[0167] $[\alpha]_D^{25+9.6^\circ}$ (c0.5, CHCl_3)

Example 9: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2,5-dimethylphenyl]-N'-[(1R)-1-(4-fluorophenyl)ethyl]urea

[0168] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2,5-dimethoxyaniline (30 mg) was added to chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1R)-1-(4-fluorophenyl)ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (35 mg, yield 77%).

[0169] $^1\text{H-NMR}$ (CDCl_3): 1.49 (3H, d, $J = 6.34$ Hz), 2.09 (3H, s), 2.13 (3H, s), 4.04 (3H, s), 4.05 (3H, s), 4.95 - 5.00 (2H, m), 6.04 (1H, s), 6.28 (1H, d, $J = 5.37$ Hz), 6.92 (1H, s), 7.04 (2H, dd, $J = 8.78, 8.78$ Hz), 7.33 (2H, dd, $J = 8.78, 5.12$ Hz), 7.42 (1H, s), 7.51 (1H, s), 7.57 (1H, s), 8.44 (1H, d, $J = 5.12$ Hz)

[0170] Mass spectrometric value (ESI-MS, m/z): 490 ($M+1$)

[0171] $[\alpha]_D^{24-11.5^\circ}$ (c0.5, CHCl_3)

Example 10: N-[3-Chloro-4-[(6,7-dimethoxy-4-quinolyl)oxy]phenyl]-N'-[1-(4-fluorophenyl)ethyl]urea

[0172] 3-Chloro-4-[(6,7-dimethoxy-4-quinolyl)oxy]aniline (83 mg) was dissolved in chloroform (2 ml) and triethylamine (0.2 ml) to prepare a solution. A solution of triphosgene (70 mg) in chloroform (0.4 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(4-fluorophenyl)ethylamine (70 mg) in chloroform (0.4 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (36 mg, yield 29%).

[0173] $^1\text{H-NMR}$ (CDCl_3): 1.48 (3H, d, $J = 6.83$ Hz), 4.01 (3H, s), 4.05 (3H, s), 4.98 (1H, m), 5.44 (1H, d, $J = 7.07$ Hz), 6.28 (1H, d, $J = 5.37$ Hz), 7.01 (2H, dd, $J = 8.78, 8.78$ Hz), 7.10 - 7.31 (4H, overlapped), 7.39 (1H, s), 7.59 (1H, s), 7.60 (1H, d, $J = 2.44$ Hz), 8.43 (1H, d, $J = 5.37$ Hz)

[0174] Mass spectrometric value (ESI-MS, m/z): 494 ($M-1$)

Example 11: N-[3-Chloro-4-[(6,7-dimethoxy-4-quinolyl)oxy]phenyl]-N'-[(1S)-1-(4-fluorophenyl)ethyl]urea

[0175] 3-Chloro-4-[(6,7-dimethoxy-4-quinolyl)oxy]aniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1S)-1-(4-fluorophenyl)ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (26 mg, yield 58%).

[0176] $^1\text{H-NMR}$ (CDCl_3): 1.48 (3H, d, $J = 6.83$ Hz), 4.01 (3H, s), 4.05 (3H, s), 4.94 - 4.99 (1H, m), 5.43 (1H, d, $J = 6.83$ Hz), 6.28 (1H, d, $J = 5.37$ Hz), 7.02 (2H, dd, $J = 8.78, 8.78$ Hz), 7.11 (1H, d, $J = 8.78$ Hz), 7.18 (s, 1H), 7.24 (1H, dd, $J = 8.78, 2.68$ Hz), 7.28 (1H, dd, $J = 5.37, 8.78$ Hz), 7.39 (1H, s), 7.59 (1H, s), 7.60 (1H, d, $J = 2.44$ Hz), 8.43 (1H, d, $J = 5.12$ Hz)

[0177] Mass spectrometric value (ESI-MS, m/z): 494 ($M-1$)

[0178] $[\alpha]_D^{24+45.1^\circ}$ (c0.5, CHCl_3)

Example 12: N-[3-Chloro-4-[(6,7-dimethoxy-4-quinolyl)oxy]phenyl]-N'-[(1R)-1-(4-fluorophenyl)ethyl]urea

[0179] 3-Chloro-4-[(6,7-dimethoxy-4-quinolyl)oxy]aniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1R)-1-(4-fluorophenyl)ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr.

The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (26 mg, yield 57%).

[0180] $^1\text{H-NMR}$ (CDCl_3): 1.50 (3H, d, $J = 6.83$ Hz), 4.03 (3H, s), 4.05 (3H, s), 4.92 - 5.00 (1H, m), 5.21 (1H, d, $J = 7.08$ Hz), 6.29 (1H, d, $J = 5.12$ Hz), 6.84 (s, 1H), 7.04 (2H, dd, $J = 8.78, 8.78$ Hz), 7.12 (1H, d, $J = 8.78$ Hz), 7.25 (1H, dd, $J = 8.78, 2.44$ Hz), 7.31 (2H, dd, $J = 8.54, 5.37$ Hz), 7.41 (1H, s), 7.59 (1H, s), 7.61 (1H, d, $J = 2.44$ Hz), 8.44 (1H, d, $J = 5.37$ Hz)

[0181] Mass spectrometric value (ESI-MS, m/z): 496 ($M+1$)

[0182] $[\alpha]_D^{25} +44.2^\circ$ (c0.5, CHCl_3)

Example 13: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methylphenyl]-N'-[1-(4-fluorophenyl)ethyl]urea

[0183] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methylaniline (78 mg) was dissolved in chloroform (2 ml) and triethylamine (0.2 ml) to prepare a solution. A solution of triphosgene (70 mg) in chloroform (0.4 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(4-fluorophenyl)ethylamine (70 mg) in chloroform (0.4 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (79 mg, yield 66%).

[0184] $^1\text{H-NMR}$ (CDCl_3): 1.49 (3H, d, $J = 6.59$ Hz), 2.13 (3H, s), 4.03 (3H, s), 4.04 (3H, s), 4.97 (1H, m), 5.05 (1H, d, $J = 6.83$ Hz), 6.14 (1H, s), 6.46 (1H, d, $J = 5.37$ Hz), 7.01 - 7.06 (4H, overlapped), 7.32 (2H, dd, $J = 8.54, 5.12$ Hz), 7.41 (1H, s), 7.52 (1H, s), 7.61 (1H, d, $J = 8.29$ Hz), 8.48 (1H, d, $J = 5.37$ Hz)

[0185] Mass spectrometric value (ESI-MS, m/z): 474 ($M-1$)

Example 14: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methylphenyl]-N'-[(1S)-1-(4-fluorophenyl)ethyl]urea

[0186] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methylaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1S)-1-(4-fluorophenyl)ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (36 mg, yield 77%).

[0187] $^1\text{H-NMR}$ (CDCl_3): 1.49 (3H, d, $J = 6.83$ Hz), 2.17 (3H, s), 4.03 (3H, s), 4.05 (3H, s), 4.87 (1H, d, $J = 6.83$ Hz), 4.94 - 4.99 (1H, m), 5.97 (1H, s), 6.47 (1H, d, $J = 5.12$ Hz), 7.01 - 7.07 (4H, m), 7.33 (2H, dd, $J = 8.54, 5.12$ Hz), 7.42 (1H, s), 7.52 (1H, s), 7.58 (1H, d, $J = 8.29$ Hz), 8.48 (1H, d, $J = 5.37$ Hz)

[0188] Mass spectrometric value (ESI-MS, m/z): 476 ($M+1$)

[0189] $[\alpha]_D^{24} +12.6^\circ$ (c0.5, CHCl_3)

Example 15: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methylphenyl]-N'-[(1R)-1-(4-fluorophenyl)ethyl]urea

[0190] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methylaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1R)-1-(4-fluorophenyl)ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (26 mg, yield 58%).

[0191] $^1\text{H-NMR}$ (CDCl_3): 1.49 (3H, d, $J = 6.59$ Hz), 2.14 (3H, s), 4.03 (3H, s), 4.04 (3H, s), 4.92 - 4.99 (1H, m), 6.03 (1H, s), 6.47 (1H, d, $J = 5.37$ Hz), 7.01 - 7.06 (4H, m), 7.33 (2H, dd, $J = 8.78, 5.37$ Hz), 7.42 (1H, s), 7.52 (1H, s), 7.59 (1H, d, $J = 9.27$ Hz), 8.48 (1H, d, $J = 5.37$ Hz)

[0192] Mass spectrometric value (ESI-MS, m/z): 476 ($M+1$)

[0193] $[\alpha]_D^{24} -14.3^\circ$ (c0.5, CHCl_3)

Example 16: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methoxyphenyl]-N'-[1-(4-fluorophenyl)ethyl]urea

[0194] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methoxyaniline (82 mg) was dissolved in chloroform (2 ml) and triethylamine (0.2 ml) to prepare a solution. A solution of triphosgene (70 mg) in chloroform (0.4 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(4-fluorophenyl)ethylamine (70 mg) in chloroform (0.4 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (58 mg, yield 47%).

[0195] ¹H-NMR (CDCl₃): 1.52 (3H, d, J = 6.83 Hz), 3.73 (3H, s), 4.03 (3H, s), 4.04 (3H, s), 4.96 (1H, m), 5.15 (1H, d, J = 6.59 Hz), 6.44 (1H, d, J = 5.12 Hz), 6.66 (1H, d, J = 2.68 Hz), 6.77 (1H, dd, J = 8.78, 2.68 Hz), 6.87 (1H, s), 7.04 (2H, dd, J = 8.78, 8.78 Hz), 7.36 (2H, dd, J = 8.72, 5.37 Hz), 7.41 (1H, s), 7.55 (1H, s), 8.16 (1H, d, J = 8.78 Hz), 8.46 (1H, d, J = 5.12 Hz)

[0196] Mass spectrometric value (ESI-MS, m/z): 490 (M-1)

Example 17: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methoxyphenyl]-N'-[(1S)-1-(4-fluorophenyl)ethyl]urea

[0197] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methoxyaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1S)-1-(4-fluorophenyl)ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (44 mg, yield 98%).

[0198] ¹H-NMR (CDCl₃): 1.52 (3H, d, J = 6.83 Hz), 3.73 (3H, s), 4.03 (3H, s), 4.04 (3H, s), 4.96 (1H, m), 5.15 (1H, d, J = 6.59 Hz), 6.44 (1H, d, J = 5.12 Hz), 6.66 (1H, d, J = 2.68 Hz), 6.77 (1H, dd, J = 8.78, 2.68 Hz), 6.87 (1H, s), 7.04 (2H, dd, J = 8.78, 8.78 Hz), 7.36 (2H, dd, J = 8.72, 5.37 Hz), 7.41 (1H, s), 7.55 (1H, s), 8.16 (1H, d, J = 8.78 Hz), 8.46 (1H, d, J = 5.12 Hz)

[0199] Mass spectrometric value (ESI-MS, m/z): 492 (M⁺+1)

[0200] [α]_D²⁴-31.6° (c0.5, CHCl₃)

Example 18: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methoxyphenyl]-N'-[(1R)-1-(4-fluorophenyl)ethyl]urea

[0201] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methoxyaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1R)-1-(4-fluorophenyl)ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (43 mg, yield 95%).

[0202] ¹H-NMR (CDCl₃): 1.53 (3H, d, J = 6.34 Hz), 3.76 (3H, s), 4.04 (3H, s), 4.04 (3H, s), 4.94 - 4.98 (2H, m), 6.44 (1H, d, J = 5.12 Hz), 6.67 (1H, d, J = 2.44 Hz), 6.76 (1H, s), 6.78 (1H, d, J = 2.44 Hz), 7.06 (2H, dd, J = 8.78, 8.78 Hz), 7.37 (2H, dd, J = 8.78, 5.37 Hz), 7.41 (1H, s), 7.55 (1H, s), 8.15 (1H, d, J = 8.78 Hz), 8.46 (1H, d, J = 5.37 Hz)

[0203] Mass spectrometric value (ESI-MS, m/z): 492 (M⁺+1)

[0204] [α]_D²⁵+30.0° (c0.5, CHCl₃)

Example 19: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-(trifluoromethyl)phenyl]-N'-[1-(4-fluorophenyl)ethyl]urea

[0205] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-(trifluoromethyl)aniline (91 mg) was dissolved in chloroform (2 ml) and triethylamine (0.2 ml) to prepare a solution. A solution of triphosgene (70 mg) in chloroform (0.4 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(4-fluorophenyl)ethylamine (70 mg) in chloroform (0.4 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (64 mg, yield 49%).

[0206] ¹H-NMR (CDCl₃): 1.52 (3H, d, J = 6.83 Hz), 4.03 (3H, s), 4.04 (3H, s), 4.94 (1H, m), 5.23 (1H, d, J = 6.59 Hz), 6.44 (1H, d, J = 5.37 Hz), 6.57 (1H, s), 7.05 (2H, dd, J = 8.78, 8.78 Hz), 7.34 (3H, overlapped), 7.39 (1H, d, J = 2.93 Hz), 7.43 (1H, s), 7.49 (1H, s), 8.14 (1H, d, J = 9.03 Hz), 8.50 (1H, d, J = 5.37 Hz)

[0207] Mass spectrometric value (ESI-MS, m/z): 528 (M-1)

Example 20: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-(trifluoromethyl)phenyl]-N'-[(1S)-1-(4-fluorophenyl)ethyl]urea

[0208] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-(trifluoromethyl)aniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1S)-1-(4-fluorophenyl)ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (9 mg, yield 20%).

[0209] ¹H-NMR (CDCl₃): 1.53 (3H, d, J = 6.83 Hz), 4.04 (3H, s), 4.05 (3H, s), 4.89 - 4.96 (1H, m), 5.07 (1H, s), 6.44 (1H, d, J = 5.37 Hz), 6.50 (1H, s), 7.06 (2H, dd, J = 8.78, 8.78 Hz), 7.33 - 7.40 (4H, m), 7.43 (1H, s), 7.49 (1H, s), 8.13

(1H, d, J = 9.03 Hz), 8.50 (1H, d, J = 5.12 Hz)

[0210] Mass spectrometric value (ESI-MS, m/z): 530 (M+1)

[0211] $[\alpha]_D^{24}$ -40.4° (c0.25, CHCl₃)

5 Example 21: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-(trifluoromethyl)phenyl]-N'-[(1R)-1-(4-fluorophenyl)ethyl]urea

[0212] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-(trifluoromethyl)aniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1R)-1-(4-fluorophenyl)ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (15 mg, yield 34%).

[0213] ¹H-NMR (CDCl₃): 1.53 (3H, d, J = 6.83 Hz), 4.04 (3H, s), 4.05 (3H, s), 4.90 - 4.96 (1H, m), 5.14 (1H, d, J = 6.59 Hz), 6.45 (1H, d, J = 5.37 Hz), 6.53 (1H, s), 7.05 (2H, dd, J = 8.78, 8.78 Hz), 7.31 - 7.36 (3H, m), 7.39 (1H, d, J = 2.68 Hz), 7.43 (1H, s), 7.49 (1H, s), 8.13 (1H, d, J = 9.03 Hz), 8.50 (1H, d, J = 5.37 Hz)

[0214] Mass spectrometric value (ESI-MS, m/z): 530 (M+1)

[0215] $[\alpha]_D^{25}$ +39.4° (c0.25, CHCl₃)

20 Example 22: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-3-methoxyphenyl]-N'-[1-(4-fluorophenyl)ethyl]urea

[0216] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-3-methoxyaniline (82 mg) was dissolved in chloroform (2 ml) and triethylamine (0.2 ml) to prepare a solution. A solution of triphosgene (70 mg) in chloroform (0.4 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(4-fluorophenyl)ethylamine (70 mg) in chloroform (0.4 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (60 mg, yield 49%).

[0217] ¹H-NMR (CDCl₃): 1.48 (3H, d, J = 6.83 Hz), 2.12 (3H, s), 2.16 (3H, s), 4.05 (3H, s), 4.05 (3H, s), 4.77 (1H, d, J = 7.07 Hz), 4.97 - 5.00 (1H, m), 5.97 (1H, s), 6.25 (1H, d, J = 5.12 Hz), 6.99 (1H, dd, J = 8.54 Hz), 7.04 (2H, dd, J = 8.78, 8.78 Hz), 7.31 (2H, dd, J = 5.12, 8.78 Hz), 7.43 (1H, s), 7.59 (1H, s), 8.44 (1H, d, J = 5.37 Hz)

[0218] Mass spectrometric value (ESI-MS, m/z): 490 (M-1)

30 Example 23: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-3-methoxyphenyl]-N'-[(1S)-1-(4-fluorophenyl)ethyl]urea

[0219] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-3-methoxyaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1S)-1-(4-fluorophenyl)ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (45 mg, yield 100%).

[0220] ¹H-NMR (CDCl₃): 1.48 (3H, d, J = 6.83 Hz), 3.71 (3H, s), 4.00 (3H, s), 4.04 (3H, s), 5.00 (1H, m), 5.48 (1H, d, J = 7.32 Hz), 6.29 (1H, d, J = 5.37 Hz), 6.62 (1H, dd, J = 8.54, 2.44 Hz), 6.99 (3H, overlapped), 7.19 (1H, s), 7.29 (1H, dd, J = 8.54, 5.12 Hz), 7.38 (1H, s), 7.47 (1H, d, J = 2.44 Hz), 7.61 (1H, s), 8.41 (1H, d, J = 5.37 Hz)

[0221] Mass spectrometric value (ESI-MS, m/z): 492 (M+1)

[0222] $[\alpha]_D^{25}$ -35.5 (c0.5, CHCl₃)

45 Example 24: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-3-methoxyphenyl]-N'-[(1R)-1-(4-fluorophenyl)ethyl]urea

[0223] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-3-methoxyaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1R)-1-(4-fluorophenyl)ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (44 mg, yield 97%).

[0224] ¹H-NMR (CDCl₃): 1.48 (3H, d, J = 6.83 Hz), 3.71 (3H, s), 4.00 (3H, s), 4.04 (3H, s), 5.00 (1H, m), 5.48 (1H, d, J = 7.32 Hz), 6.29 (1H, d, J = 5.37 Hz), 6.62 (1H, dd, J = 8.54, 2.44 Hz), 6.99 (3H, overlapped), 7.19 (1H, s), 7.29 (1H, dd, J = 8.54, 5.12 Hz), 7.38 (1H, s), 7.47 (1H, d, J = 2.44 Hz), 7.61 (1H, s), 8.41 (1H, d, J = 5.37 Hz)

[0225] Mass spectrometric value (ESI-MS, m/z): 492 (M+1)

[0226] $[\alpha]_D^{25}$ +35.8° (c0.5, CHCl₃)

Example 25: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2,3-dimethylphenyl]-N'-[1-(4-fluorophenyl)ethyl]urea

[0227] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2,3-dimethylaniline (81 mg) was dissolved in chloroform (2 ml) and triethylamine (0.2 ml) to prepare a solution. A solution of triphosgene (70 mg) in chloroform (0.4 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(4-fluorophenyl)ethylamine (70 mg) in chloroform (0.4 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (6 mg, yield 5%).

[0228] ¹H-NMR (CDCl₃): 1.48 (3H, d, J = 6.83 Hz), 2.11 (3H, s), 2.16 (3H, s), 4.06 (6H, s), 4.78 (1H, d, J = 5.86 Hz), 5.00 (1H, m), 5.97 (1H, s), 6.26 (1H, d, J = 5.37 Hz), 6.90 - 7.25 (5H, overlapped), 7.30 (2H, m), 7.46 (1H, s), 7.59 (1H, s), 8.44 (1H, d, J = 5.37 Hz)

[0229] Mass spectrometric value (ESI-MS, m/z): 488 (M-1)

Example 26: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2,3-dimethylphenyl]-N'-[(1S)-1-(4-fluorophenyl)ethyl]urea

[0230] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2,3-dimethylaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. A solution of (1S)-1-(4-fluorophenyl)ethylamine (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (34 mg, yield 76%).

[0231] ¹H-NMR (CDCl₃): 1.48 (3H, d, J = 6.83 Hz), 2.11 (3H, s), 2.16 (3H, s), 4.06 (6H, s), 4.78 (1H, d, J = 5.86 Hz), 5.00 (1H, m), 5.97 (1H, s), 6.26 (1H, d, J = 5.37 Hz), 6.90 - 7.25 (5H, overlapped), 7.30 (2H, m), 7.46 (1H, s), 7.59 (1H, s), 8.44 (1H, d, J = 5.37 Hz)

[0232] Mass spectrometric value (ESI-MS, m/z): 488 (M-1)

[0233] [α]_D²⁵+30.0° (c0.5, CHCl₃)

Example 27: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2,3-dimethylphenyl]-N'-[(1R)-1-(4-fluorophenyl)ethyl]urea

[0234] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2,3-dimethylaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1R)-1-(4-fluorophenyl)ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (37 mg, yield 81%).

[0235] ¹H-NMR (CDCl₃): 1.48 (3H, d, J = 6.83 Hz), 2.12 (3H, s), 2.16 (3H, s), 4.05 (3H, s), 4.05 (3H, s), 4.77 (1H, d, J = 7.32 Hz), 4.98 - 5.04 (1H, m), 5.99 (1H, s), 6.25 (1H, d, J = 5.37 Hz), 6.99 (1H, d, J = 8.78 Hz), 7.04 (2H, dd, J = 8.78, 8.78 Hz), 7.31 (2H, dd, J = 5.12, 8.78 Hz), 7.43 (1H, s), 7.59 (1H, s), 8.44 (1H, d, J = 5.12 Hz)

[0236] Mass spectrometric value (ESI-MS, m/z): 488 (M-1)

[0237] [α]_D²⁴-35.6° (c0.5, CHCl₃)

Example 28: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-fluorophenyl]-N'-[1-(4-fluorophenyl)ethyl]urea

[0238] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-fluoroaniline (79 mg) was dissolved in chloroform (2 ml) and triethylamine (0.2 ml) to prepare a solution. A solution of triphosgene (70 mg) in chloroform (0.4 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(4-fluorophenyl)ethylamine (70 mg) in chloroform (0.4 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (39 mg, yield 33%).

[0239] ¹H-NMR (CDCl₃): 1.50 (3H, d, J = 6.83 Hz), 4.03 (6H, s), 4.97 (1H, m), 5.55 (1H, d, J = 6.59 Hz), 6.46 (1H, d, J = 5.37 Hz), 6.87 - 6.96 (3H, overlapped), 7.01 (1H, dd, J = 8.78, 8.78 Hz), 7.33 (2H, dd, J = 8.54, 5.37 Hz), 7.42 (1H, s), 7.50 (1H, s), 8.17 (1H, dd, J = 9.03, 9.03 Hz), 8.48 (1H, d, J = 5.37 Hz)

[0240] Mass spectrometric value (ESI-MS, m/z): 478 (M-1)

Example 29: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-fluorophenyl]-N'-[(1S)-1-(4-fluorophenyl)ethyl]urea

[0241] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-fluoroaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution,

and the mixture was stirred at room temperature for one hr. Next, a solution of (1S)-1-(4-fluorophenyl)ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (35 mg, yield 76%).

[0242] ¹H-NMR (CDCl₃): 1.50 (3H, d, J = 6.83 Hz), 4.03 (6H, s), 4.97 (1H, m), 5.55 (1H, d, J = 6.59 Hz), 6.46 (1H, d, J = 5.37 Hz), 6.87 - 6.96 (3H, overlapped), 7.01 (1H, dd, J = 8.78, 8.78 Hz), 7.33 (2H, dd, J = 8.54, 5.37 Hz), 7.42 (1H, s), 7.50 (1H, s), 8.17 (1H, dd, J = 9.03, 9.03 Hz), 8.48 (1H, d, J = 5.37 Hz)

[0243] Mass spectrometric value (ESI-MS, m/z): 480 (M⁺+1)

[0244] [α]_D²⁴-45.8° (c0.5, CHCl₃)

Example 30: N-{4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-fluorophenyl}-N'-[(1R)-1-(4-fluorophenyl)ethyl]urea

[0245] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-fluoroaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1R)-1-(4-fluorophenyl)ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (34 mg, yield 74%).

[0246] ¹H-NMR (CDCl₃): 1.50 (3H, d, J = 6.83 Hz), 4.03 (6H, s), 4.97 (1H, m), 5.55 (1H, d, J = 6.59 Hz), 6.46 (1H, d, J = 5.37 Hz), 6.87 - 6.96 (3H, overlapped), 7.01 (1H, dd, J = 8.78, 8.78 Hz), 7.33 (2H, dd, J = 8.54, 5.37 Hz), 7.42 (1H, s), 7.50 (1H, s), 8.17 (1H, dd, J = 9.03, 9.03 Hz), 8.48 (1H, d, J = 5.37 Hz)

[0247] Mass spectrometric value (ESI-MS, m/z): 480 (M⁺+1)

[0248] [α]_D²⁵+45.0° (c0.5, CHCl₃)

Example 31: N-{4-[(6,7-Dimethoxy-4-quinolyl)oxy]-3-fluorophenyl}-N'-[1-(4-fluorophenyl)ethyl]urea

[0249] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-3-fluoroaniline (79 mg) was dissolved in chloroform (2 ml) and triethylamine (0.2 ml) to prepare a solution. A solution of triphosgene (70 mg) in chloroform (0.4 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(4-fluorophenyl)ethylamine (70 mg) in chloroform (0.4 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (55 mg, yield 46%).

[0250] ¹H-NMR (CDCl₃): 1.48 (3H, d, J = 6.83 Hz), 4.00 (3H, s), 4.05 (3H, s), 4.98 (1H, m), 5.55 (1H, d, J = 7.08 Hz), 6.37 (1H, d, J = 5.37 Hz), 7.00 (2H, dd, J = 8.54, 8.54 Hz), 7.00 (1H, overlapped), 7.10 (1H, dd, J = 8.78, 8.78 Hz), 7.28 (2H, dd, J = 8.54, 5.37 Hz), 7.33 (1H, s), 7.38 (1H, s), 7.46 (1H, dd, J = 12.18, 2.44 Hz), 7.58 (1H, s), 8.44 (1H, d, J = 5.37 Hz)

[0251] Mass spectrometric value (ESI-MS, m/z): 478 (M-1)

Example 32: N-{4-[(6,7-Dimethoxy-4-quinolyl)oxy]-3-fluorophenyl}-N'-[(1S)-1-(4-fluorophenyl)ethyl]urea

[0252] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-3-fluoroaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1S)-1-(4-fluorophenyl)ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (27 mg, yield 59%).

[0253] ¹H-NMR (CDCl₃): 1.48 (3H, d, J = 6.83 Hz), 4.00 (3H, s), 4.05 (3H, s), 4.98 (1H, m), 5.55 (1H, d, J = 7.08 Hz), 6.37 (1H, d, J = 5.37 Hz), 7.00 (2H, dd, J = 8.54, 8.54 Hz), 7.00 (1H, overlapped), 7.10 (1H, dd, J = 8.78, 8.78 Hz), 7.28 (2H, dd, J = 8.54, 5.37 Hz), 7.33 (1H, s), 7.38 (1H, s), 7.46 (1H, dd, J = 12.18, 2.44 Hz), 7.58 (1H, s), 8.44 (1H, d, J = 5.37 Hz)

[0254] Mass spectrometric value (ESI-MS, m/z): 480 (M⁺+1)

[0255] [α]_D²⁴-45.4° (c0.5, CHCl₃)

Example 33: N-{4-[(6,7-Dimethoxy-4-quinolyl)oxy]-3-fluorophenyl}-N'-[(1R)-1-(4-fluorophenyl)ethyl]urea

[0256] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-3-fluoroaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1R)-1-(4-fluorophenyl)ethylamine (35

mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (30 mg, yield 66%).

[0257] $^1\text{H-NMR}$ (CDCl_3): 1.48 (3H, d, $J = 6.83$ Hz), 4.00 (3H, s), 4.05 (3H, s), 4.98 (1H, m), 5.55 (1H, d, $J = 7.08$ Hz), 6.37 (1H, d, $J = 5.37$ Hz), 7.00 (2H, dd, $J = 8.54, 8.54$ Hz), 7.00 (1H, overlapped), 7.10 (1H, dd, $J = 8.78, 8.78$ Hz), 7.28 (2H, dd, $J = 8.54, 5.37$ Hz), 7.33 (1H, s), 7.38 (1H, s), 7.46 (1H, dd, $J = 12.18, 2.44$ Hz), 7.58 (1H, s), 8.44 (1H, d, $J = 5.37$ Hz)

[0258] Mass spectrometric value (ESI-MS, m/z): 480 ($M^+ + 1$)

[0259] $[\alpha]_D^{25} + 44.7^\circ$ (c0.5, CHCl_3)

Example 34: N-[(1S)-1-(4-Bromophenyl)ethyl]-N'-[4-[(6,7-dimethoxy-4-quinolyl)oxy]phenyl]urea

[0260] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]aniline (70 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (33 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for 75 min. Next, a solution of (1S)-1-(4-bromophenyl)ethylamine (46 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (68 mg, yield 55%).

[0261] $^1\text{H-NMR}$ (CDCl_3): 1.45 (3H, d, $J = 7.07$ Hz), 4.02 (3H, s), 4.04 (3H, s), 4.95 (1H, m), 5.37 (1H, d, $J = 7.32$ Hz), 6.42 (1H, d, $J = 5.36$ Hz), 7.00 (1H, s), 7.08 (2H, dd, $J = 8.78$ Hz), 7.20 (2H, d, $J = 8.29$ Hz), 7.37 (2H, dd, $J = 8.78$ Hz), 7.39 (1H, s), 7.44 (2H, d, $J = 8.29$ Hz), 7.55 (1H, s), 8.45 (1H, d, $J = 5.36$ Hz)

[0262] Mass spectrometric value (ESI-MS, m/z): 520, 522 ($M-1$)

[0263] $[\alpha]_D^{25} - 33.0^\circ$ (c0.5, CHCl_3)

Example 35: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]phenyl]-N'-[(1S)-1-(4-nitrophenyl)ethyl]urea

[0264] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]aniline (70 mg) was dissolved in chloroform (1 ml) and triethylamine (0.14 ml) to prepare a solution. A solution of triphosgene (33 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for 75 min. Next, a solution of (1S)-1-(4-nitrophenyl)ethylamine hydrochloride (47 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (69 mg, yield 60%).

[0265] $^1\text{H-NMR}$ (CDCl_3): 1.52 (3H, d, $J = 7.07$ Hz), 4.03 (3H, s), 4.04 (3H, s), 5.11 (1H, m), 5.26 (1H, d, $J = 6.83$ Hz), 6.43 (1H, d, $J = 5.36$ Hz), 6.81 (1H, s), 7.12 (2H, d, $J = 8.78$ Hz), 7.40 (2H, d, $J = 8.78$ Hz), 7.41 (1H, s), 7.51 (2H, d, $J = 8.78$ Hz), 7.55 (1H, s), 8.19 (2H, d, $J = 8.78$ Hz), 8.46 (1H, d, $J = 5.36$ Hz)

[0266] Mass spectrometric value (ESI-MS, m/z): 487 ($M-1$)

[0267] $[\alpha]_D^{25} - 27.0^\circ$ (c0.25, CHCl_3)

Example 36: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]phenyl]-N'-[(1R)-1-(4-nitrophenyl)ethyl]urea

[0268] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]aniline (70 mg) was dissolved in chloroform (1 ml) and triethylamine (0.14 ml) to prepare a solution. A solution of triphosgene (33 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for 75 min. Next, a solution of (1R)-1-(4-nitrophenyl)ethylamine hydrochloride (47 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (16 mg, yield 13%).

[0269] $^1\text{H-NMR}$ (CDCl_3): 1.52 (3H, d, $J = 7.07$ Hz), 4.03 (3H, s), 4.04 (3H, s), 5.11 (1H, m), 5.24 (1H, d, $J = 6.83$ Hz), 6.43 (1H, d, $J = 5.36$ Hz), 6.76 (1H, s), 7.12 (2H, d, $J = 8.78$ Hz), 7.40 (2H, d, $J = 8.78$ Hz), 7.41 (1H, s), 7.51 (2H, d, $J = 8.78$ Hz), 7.55 (1H, s), 8.19 (2H, d, $J = 8.78$ Hz), 8.46 (1H, d, $J = 5.36$ Hz)

[0270] Mass spectrometric value (ESI-MS, m/z): 487 ($M-1$)

[0271] $[\alpha]_D^{25} + 28.0^\circ$ (c0.5, CHCl_3)

Example 37: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]phenyl]-N'-[(1S)-1-phenylpropyl]urea

[0272] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]aniline (70 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (33 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for 75 min. Next, a solution of (1S)-1-phenylpropylamine (31 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was

purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (62 mg, yield 57%).

[0273] $^1\text{H-NMR}$ (CDCl_3): 0.91 (3H, d, $J = 7.56$ Hz), 1.82 (2H, m), 4.02 (3H, s), 4.03 (3H, s), 4.71 (1H, m), 5.32 (1H, d, $J = 7.08$ Hz), 6.41 (1H, d, $J = 5.37$ Hz), 6.87 (1H, s), 7.07 (2H, d, $J = 9.02$ Hz), 7.28 - 7.36 (8H, overlapped), 7.40 (1H, s), 7.54 (1H, s), 8.45 (1H, d, $J = 5.37$ Hz)

[0274] Mass spectrometric value (ESI-MS, m/z): 456 (M-1)

[0275] $[\alpha]_D^{25}$ -34.7° (c0.25, CHCl_3)

Example 38: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]phenyl]-N'-[(1R)-1-phenylpropyl]urea

[0276] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]aniline (70 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (33 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for 75 min. Next, a solution of (1R)-1-phenylpropylamine (31 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (75 mg, yield 69%).

[0277] $^1\text{H-NMR}$ (CDCl_3): 0.93 (3H, d, $J = 7.56$ Hz), 1.84 (2H, m), 4.03 (3H, s), 4.04 (3H, s), 4.70 (1H, m), 5.09 (1H, d, $J = 7.08$ Hz), 6.42 (1H, d, $J = 5.37$ Hz), 6.54 (1H, s), 7.08 (2H, d, $J = 9.03$ Hz), 7.29 - 7.38 (8H, overlapped), 7.41 (1H, s), 7.54 (1H, s), 8.45 (1H, d, $J = 5.37$ Hz)

[0278] Mass spectrometric value (ESI-MS, m/z): 456 (M-1)

[0279] $[\alpha]_D^{25}$ +34.4° (c0.5, CHCl_3)

Example 41: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]phenyl]-N'-[(1S)-1-(4-methylphenyl)ethyl]urea

[0280] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]aniline (70 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (33 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for 75 min. Next, a solution of (1S)-1-(4-methylphenyl)ethylamine (31 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (52 mg, yield 48%).

[0281] $^1\text{H-NMR}$ (CDCl_3): 1.49 (3H, d, $J = 6.83$ Hz), 2.33 (3H, s), 4.02 (3H, s), 4.03 (3H, s), 4.53 (1H, m), 5.15 (1H, d, $J = 6.83$ Hz), 6.41 (1H, d, $J = 5.37$ Hz), 6.68 (1H, s), 7.07 (2H, d, $J = 8.78$ Hz), 7.15 (2H, d, $J = 7.81$ Hz), 7.24 (2H, d, $J = 7.81$ Hz), 7.35 (2H, d, $J = 8.78$ Hz), 7.40 (1H, s), 7.54 (1H, s), 8.45 (1H, d, $J = 5.37$ Hz)

[0282] Mass spectrometric value (ESI-MS, m/z): 456 (M-1)

[0283] $[\alpha]_D^{24}$ -32.0° (c0.5, CHCl_3)

Example 42: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]phenyl]-N'-[(1R)-1-(4-methylphenyl)ethyl]urea

[0284] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]aniline (70 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (33 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for 75 min. Next, a solution of (1R)-1-(4-methylphenyl)ethylamine (31 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (68 mg, yield 63%).

[0285] $^1\text{H-NMR}$ (CDCl_3): 1.49 (3H, d, $J = 6.83$ Hz), 2.33 (3H, s), 4.02 (3H, s), 4.03 (3H, s), 4.93 (1H, m), 5.18 (1H, d, $J = 7.07$ Hz), 6.41 (1H, d, $J = 5.37$ Hz), 6.72 (1H, s), 7.07 (2H, d, $J = 8.78$ Hz), 7.15 (2H, d, $J = 8.05$ Hz), 7.24 (2H, d, $J = 8.05$ Hz), 7.35 (2H, d, $J = 8.78$ Hz), 7.40 (1H, s), 7.54 (1H, s), 8.45 (1H, d, $J = 5.37$ Hz)

[0286] Mass spectrometric value (ESI-MS, m/z): 456 (M-1)

[0287] $[\alpha]_D^{24}$ +31.7° (c0.5, CHCl_3)

Example 43: N-[(1S)-2,3-Dihydro-1H-1-indenyl]-N'-[4-[(6,7-dimethoxy-4-quinolyl)oxy]phenyl]urea

[0288] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]aniline (70 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (33 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for 75 min. Next, a solution of (S)-1-aminoindane (31 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (43 mg, yield 40%).

[0289] $^1\text{H-NMR}$ ($\text{CDCl}_3+\text{CD}_3\text{OD}$): 1.83 (1H, m), 2.64 (1H, m), 2.89 (1H, m), 2.99 (1H, m), 4.06 (3H, s), 4.07 (3H, s), 5.38 (1H, t, $J = 7.32$ Hz), 6.47 (1H, d, $J = 5.37$ Hz), 7.11 (2H, d, $J = 8.54$ Hz), 7.23 (3H, overlapped), 7.37 (2H, overlapped), 7.52 (2H, d, $J = 8.54$ Hz), 7.59 (1H, s), 8.40 (1H, d, $J = 5.37$ Hz)

[0290] Mass spectrometric value (ESI-MS, m/z): 454 (M-1)

[0291] $[\alpha]_D^{24} + 10.0^\circ$ (c0.5, CHCl_3)

Example 44: N-[(1R)-2,3-Dihydro-1H-1-indenyl]-N'-[4-[(6,7-dimethoxy-4-quinolyl)oxy]phenyl]urea

[0292] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]aniline (70 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (33 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for 75 min. Next, a solution of (R)-1-aminoindane (31 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (35 mg, yield 32%).

[0293] $^1\text{H-NMR}$ (CDCl_3): 1.81 (1H, m), 2.63 (1H, m), 2.87 (1H, m), 2.97 (1H, m), 4.05 (6H, s), 5.39 (1H, t, $J = 7.56$ Hz), 6.44 (1H, d, $J = 5.37$ Hz), 7.10 (2H, d, $J = 8.78$ Hz), 7.23 (3H, overlapped), 7.36 (1H, brd, $J = 6.83$ Hz), 7.38 (1H, s), 7.57 (1H, s), 8.41 (1H, d, $J = 5.37$ Hz)

[0294] Mass spectrometric value (ESI-MS, m/z): 454 (M-1)

[0295] $[\alpha]_D^{24-9.5^\circ}$ (c0.25, CHCl_3)

Example 45: N-[(1R)-1-(3-Bromophenyl)ethyl]-N'-[4-[(6,7-dimethoxy-4-quinolyl)oxy]phenyl]urea

[0296] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]aniline (70 mg) was dissolved in chloroform (1 ml) and triethylamine (0.14 ml) to prepare a solution. A solution of triphosgene (33 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for 75 min. Next, a solution of (1R)-1-(3-bromophenyl)ethylamine hydrochloride (54 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (70 mg, yield 57%).

[0297] $^1\text{H-NMR}$ (CDCl_3): 1.43 (3H, d, $J = 6.83$ Hz), 4.01 (3H, s), 4.03 (3H, s), 4.97 (1H, m), 5.72 (1H, d, $J = 7.32$ Hz), 6.41 (1H, d, $J = 5.37$ Hz), 7.07 (2H, d, $J = 8.78$ Hz), 7.16 (1H, t, $J = 7.81$ Hz), 7.24 (1H, d, $J = 7.81$ Hz), 7.25 - 7.40 (3H, overlapped), 7.44 (2H, overlapped), 7.55 (1H, s), 8.44 (1H, d, $J = 5.37$ Hz)

[0298] Mass spectrometric value (ESI-MS, m/z): 520, 522 (M-1)

[0299] $[\alpha]_D^{24} + 34.3^\circ$ (c0.5, CHCl_3)

Example 46: N-[(1S)-1-(3-Bromophenyl)ethyl]-N'-[4-[(6,7-dimethoxy-4-quinolyl)oxy]phenyl]urea

[0300] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]aniline (70 mg) was dissolved in chloroform (1 ml) and triethylamine (0.14 ml) to prepare a solution. A solution of triphosgene (33 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for 75 min. Next, a solution of (1S)-1-(3-bromophenyl)ethylamine hydrochloride (54 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (76 mg, yield 62%).

[0301] $^1\text{H-NMR}$ (CDCl_3): 1.46 (3H, d, $J = 6.83$ Hz), 4.02 (3H, s), 4.04 (3H, s), 4.98 (1H, m), 5.38 (1H, d, $J = 7.32$ Hz), 6.42 (1H, d, $J = 5.37$ Hz), 6.69 (1H, s), 7.09 (2H, d, $J = 9.03$ Hz), 7.19 (1H, t, $J = 7.81$ Hz), 7.26 (1H, overlapped with CHCl_3), 7.37 - 7.40 (4H, overlapped), 7.47 (1H, s), 7.55 (1H, s), 8.45 (1H, d, $J = 5.37$ Hz)

[0302] Mass spectrometric value (ESI-MS, m/z): 520, 522 (M-1)

[0303] $[\alpha]_D^{24} - 33.5^\circ$ (c0.5, CHCl_3)

Example 47: N-[(1S)-1-(4-Chlorophenyl)ethyl]-N'-[4-[(6,7-dimethoxy-4-quinolyl)oxy]phenyl]urea

[0304] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]aniline (70 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (33 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for 75 min. Next, a solution of (1S)-1-(4-chlorophenyl)ethylamine (36 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (61 mg, yield 54%).

[0305] $^1\text{H-NMR}$ ($\text{CDCl}_3+\text{CD}_3\text{OD}$): 1.47 (3H, d, $J = 6.83$ Hz), 4.05 (3H, s), 4.06 (3H, s), 4.95 (1H, q, $J = 6.83$ Hz), 6.44 (1H, d, $J = 5.37$ Hz), 7.08 (2H, d, $J = 8.78$ Hz), 7.31 (4H, singletlike, overlapped), 7.36 (1H, s), 7.47 (2H, d, $J =$

8.78 Hz), 7.58 (1H, s), 8.37 (1H, d, J = 5.37 Hz)

[0306] Mass spectrometric value (ESI-MS, m/z): 476 (M-1)

[0307] $[\alpha]_D^{24-30.3^\circ}$ (c0.5, CHCl₃)

5 Example 48: N-[(1R)-1-(4-Chlorophenyl)ethyl]-N'-[4-[(6,7-dimethoxy-4-quinolyl)oxy]phenyl]urea

[0308] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]aniline (70 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (33 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for 75 min. Next, a solution of (1R)-1-(4-chlorophenyl)ethylamine (36 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (70 mg, yield 62%).

[0309] ¹H-NMR (CDCl₃): 1.46 (3H, d, J = 6.83 Hz), 4.03 (3H, s), 4.04 (3H, s), 4.96 (1H, q, J = 6.83 Hz), 6.41 (1H, d, J = 5.37 Hz), 7.06 (2H, d, J = 9.03 Hz), 7.28 (4H, doubletlike, overlapped), 7.37 (1H, s), 7.42 (2H, d, J = 9.03 Hz), 7.56 (1H, s), 8.40 (1H, d, J = 5.37 Hz)

[0310] Mass spectrometric value (ESI-MS, m/z): 476 (M-1)

[0311] $[\alpha]_D^{24+30.3^\circ}$ (c0.5, CHCl₃)

20 Example 49: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]phenyl]-N'-[(1S)-1-(3-methoxyphenyl)ethyl]urea

[0312] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]aniline (70 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (33 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for 75 min. Next, a solution of (1S)-1-(3-methoxyphenyl)ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (67 mg, yield 60%).

[0313] ¹H-NMR (CDCl₃): 1.47 (3H, d, J = 6.83 Hz), 3.79 (3H, s), 4.04 (6H, s), 4.95 (1H, q, J = 6.83 Hz), 6.41 (1H, d, J = 5.37 Hz), 6.79 (1H, dd, J = 7.81, 2.20 Hz), 6.90 (1H, d, J = 2.20 Hz), 6.94 (1H, d, J = 7.81 Hz), 7.06 (2H, d, J = 8.78 Hz), 7.25 (1H, t, J = 7.81 Hz), 7.38 (1H, s), 7.41 (2H, d, J = 8.78 Hz), 7.56 (1H, s), 8.40 (1H, d, J = 5.37 Hz)

[0314] Mass spectrometric value (ESI-MS, m/z): 472 (M-1)

[0315] $[\alpha]_D^{24-25.2^\circ}$ (c0.5, CHCl₃)

30 Example 50: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]phenyl]-N'-[(1R)-1-(3-methoxyphenyl)ethyl]urea

[0316] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]aniline (70 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (33 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for 75 min. Next, a solution of (1R)-1-(3-methoxyphenyl)ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (53 mg, yield 47%).

[0317] ¹H-NMR (CDCl₃): 1.46 (3H, d, J = 6.83 Hz), 3.75 (3H, s), 4.00 (3H, s), 4.03 (3H, s), 4.95 (1H, m), 5.59 (1H, d, J = 7.07 Hz), 6.40 (1H, d, J = 5.37 Hz), 6.77 (1H, dd, J = 7.81, 2.20 Hz), 6.88 (1H, d, J = 2.20 Hz), 6.91 (1H, d, J = 7.81 Hz), 7.05 (2H, d, J = 8.78 Hz), 7.23 (1H, t, J = 7.81 Hz), 7.28 (1H, s), 7.36 (2H, d, J = 8.78 Hz), 7.39 (1H, s), 7.54 (1H, s), 8.43 (1H, d, J = 5.37 Hz)

[0318] Mass spectrometric value (ESI-MS, m/z): 472 (M-1)

[0319] $[\alpha]_D^{24+25.8^\circ}$ (c0.5, CHCl₃)

45 Example 51: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]phenyl]-N'-[(1S)-1-(2-methoxyphenyl)ethyl]urea

[0320] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]aniline (70 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (33 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for 75 min. Next, a solution of (1S)-1-(2-methoxyphenyl)ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (67 mg, yield 59%).

[0321] ¹H-NMR (CDCl₃): 1.49 (3H, d, J = 6.83 Hz), 3.81 (3H, s), 4.01 (3H, s), 4.04 (3H, s), 5.19 (1H, m), 5.74 (1H, br), 6.42 (1H, d, J = 5.37 Hz), 6.90 (1H, d, J = 7.81 Hz), 6.94 (1H, t, J = 6.59 Hz), 7.08 (2H, d, J = 8.78 Hz), 7.22 - 7.29 (2H, overlapped), 7.38 (2H, d, J = 8.78 Hz), 7.40 (1H, s), 7.55 (1H, s), 8.46 (1H, d, J = 5.37 Hz)

[0322] Mass spectrometric value (ESI-MS, m/z): 472 (M-1)

[0323] $[\alpha]_D^{24}$ -17.7° (c0.5, CHCl₃)

Example 52: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]phenyl]-N'-[(1R)-1-(2-methoxyphenyl)ethyl]urea

[0324] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]aniline (70 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (33 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for 75 min. Next, a solution of (1R)-1-(2-methoxyphenyl)ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (64 mg, yield 57%).

[0325] ¹H-NMR (CDCl₃): 1.48 (3H, d, J = 6.83 Hz), 3.76 (3H, s), 4.00 (3H, s), 4.04 (3H, s), 5.20 (1H, m), 5.86 (1H, br), 6.42 (1H, d, J = 5.37 Hz), 6.87 (1H, d, J = 8.29 Hz), 6.92 (1H, t, J = 7.32 Hz), 7.07 (2H, d, J = 9.03 Hz), 7.22 (1H, t, J = 7.32 Hz), 7.26 (1H, overlapped with CHCl₃), 7.39 (2H, d, J = 9.03 Hz), 7.40 (1H, s), 7.55 (1H, s), 8.43 (1H, d, J = 5.37 Hz)

[0326] Mass spectrometric value (ESI-MS, m/z): 472 (M-1)

[0327] $[\alpha]_D^{24}$ +17.8° (c0.5, CHCl₃)

Example 53: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]phenyl]-N'-[(1S)-1-(4-methoxyphenyl)ethyl]urea

[0328] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]aniline (70 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (33 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for 75 min. Next, a solution of (1S)-1-(4-methoxyphenyl)ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (63 mg, yield 57%).

[0329] ¹H-NMR (CDCl₃): 1.48 (3H, d, J = 6.83 Hz), 3.78 (3H, s), 4.02 (3H, s), 4.03 (3H, s), 4.92 (1H, m), 5.26 (1H, d, J = 6.83 Hz), 6.41 (1H, d, J = 5.37 Hz), 6.87 (2H, d, J = 8.78 Hz), 6.87 (1H, s), 7.07 (2H, d, J = 8.78 Hz), 7.27 (2H, d, J = 8.78 Hz), 7.35 (2H, d, J = 8.78 Hz), 7.39 (1H, s), 7.54 (1H, s), 8.44 (1H, d, J = 5.37 Hz)

[0330] Mass spectrometric value (ESI-MS, m/z): 472 (M-1)

[0331] $[\alpha]_D^{24}$ -33.0° (c0.5, CHCl₃)

Example 54: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]phenyl]-N'-[(1R)-1-(4-methoxyphenyl)ethyl]urea

[0332] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]aniline (70 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (33 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for 75 min. Next, a solution of (1R)-1-(4-methoxyphenyl)ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (56 mg, yield 50%).

[0333] ¹H-NMR (CDCl₃): 1.47 (3H, d, J = 6.83 Hz), 3.77 (3H, s), 4.01 (3H, s), 4.03 (3H, s), 4.92 (1H, m), 5.37 (1H, d, J = 6.83 Hz), 6.40 (1H, d, J = 5.37 Hz), 6.85 (2H, d, J = 8.72 Hz), 7.03 (1H, s), 7.06 (2H, d, J = 9.03 Hz), 7.25 (2H, d, J = 9.03 Hz), 7.35 (2H, d, J = 8.78 Hz), 7.39 (1H, s), 7.54 (1H, s), 8.44 (1H, d, J = 5.37 Hz)

[0334] Mass spectrometric value (ESI-MS, m/z): 472 (M-1)

[0335] $[\alpha]_D^{24}$ +34.3° (c0.5, CHCl₃)

Example 55: N-[(1R)-1-(3-Bromophenyl)ethyl]-N'-[2-chloro-4-[(6,7-dimethoxy-4-quinolyl)oxy]phenyl]urea

[0336] 2-Chloro-4-[(6,7-dimethoxy-4-quinolyl)oxy]aniline (83 mg) was dissolved in chloroform (2 ml) and triethylamine (0.2 ml) to prepare a solution. A solution of triphosgene (70 mg) in chloroform (0.4 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1R)-1-(3-bromophenyl)ethylamine (70 mg) in chloroform (0.4 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (64 mg, yield 46%).

[0337] ¹H-NMR (CDCl₃): 1.49 (3H, d, J = 6.08 Hz), 4.02 (3H, s), 4.03 (3H, s), 4.95 (1H, m), 5.89 (1H, d, J = 6.59 Hz), 6.46 (1H, d, J = 5.37 Hz), 7.05 - 7.10 (2H, overlapped), 7.16 (1H, d, J = 2.68 Hz), 7.20 (1H, dd, J = 7.81, 7.81 Hz), 7.30 (1H, d, J = 7.81 Hz), 7.38 (1H, m), 7.41 (1H, s), 7.50 (1H, s), 7.52 (1H, s), 8.27 (1H, d, J = 9.03 Hz), 8.48 (1H, d, J = 5.37 Hz)

[0338] Mass spectrometric value (ESI-MS, m/z): 554, 556 (M-1)

[0339] $[\alpha]_D^{24} + 32.3^\circ$ (c0.5, CHCl₃)

Example 56: N-[(1R)-1-(3-Bromophenyl)ethyl]-N'-[4-[(6,7-dimethoxy-4-quinolyl)oxy]-2,5-dimethylphenyl]urea

[0340] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2,5-dimethylaniline (83 mg) was dissolved in chloroform (2 ml) and triethylamine (0.2 ml) to prepare a solution. A solution of triphosgene (70 mg) in chloroform (0.4 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1R)-1-(3-bromophenyl)ethylamine (70 mg) in chloroform (0.4 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (63 mg, yield 45%).

[0341] ¹H-NMR (CDCl₃): 1.47 (3H, d, J = 7.08 Hz), 2.12 (3H, s), 2.13 (3H, s), 4.04 (3H, s), 4.05 (3H, s), 4.97 (1H, m), 5.26 (1H, d, J = 7.32 Hz), 6.26 (1H, s), 6.29 (1H, d, J = 5.37 Hz), 6.92 (1H, s), 7.21 (1H, dd, J = 7.81, 7.81 Hz), 7.28 (1H, d, J = 7.81 Hz), 7.39 (1H, d, J = 7.81 Hz), 7.42 (1H, s), 7.48 (1H, s), 7.54 (1H, s), 7.57 (1H, s), 8.44 (1H, d, J = 5.37 Hz)

[0342] Mass spectrometric value (ESI-MS, m/z): 548, 550 (M-1)

[0343] $[\alpha]_D^{24} - 7.6^\circ$ (c0.5, CHCl₃)

Example 57: N-[(1R)-1-(3-Bromophenyl)ethyl]-N'-[3-chloro-4-[(6,7-dimethoxy-4-quinolyl)oxy]phenyl]urea

[0344] 3-Chloro-4-[(6,7-dimethoxy-4-quinolyl)oxy]aniline (78 mg) was dissolved in chloroform (2 ml) and triethylamine (0.2 ml) to prepare a solution. A solution of triphosgene (70 mg) in chloroform (0.4 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1R)-1-(3-bromophenyl)ethylamine (70 mg) in chloroform (0.4 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (41 mg, yield 31%).

[0345] ¹H-NMR (CDCl₃): 1.45 (3H, d, J = 6.83 Hz), 4.01 (3H, s), 4.05 (3H, s), 4.95 (1H, m), 5.87 (1H, d, J = 7.32 Hz), 6.28 (1H, d, J = 5.37 Hz), 7.09 (1H, d, J = 8.78 Hz), 7.17 (1H, dd, J = 7.81, 7.81 Hz), 7.26 (2H, overlapped), 7.35 (1H, d, J = 7.81 Hz), 7.40 (1H, s), 7.46 (1H, s), 7.59 (1H, s), 7.62 (1H, d, J = 2.44 Hz), 7.67 (1H, s), 8.41 (1H, d, J = 5.37 Hz)

[0346] Mass spectrometric value (ESI-MS, m/z): 554, 556 (M-1)

[0347] $[\alpha]_D^{24} + 43.4^\circ$ (c0.5, CHCl₃)

Example 58: N-[(1R)-1-(3-Bromophenyl)ethyl]-N'-[4-[(6,7-dimethoxy-4-quinolyl)oxy]-2-methylphenyl]urea

[0348] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methylaniline (82 mg) was dissolved in chloroform (2 ml) and triethylamine (0.2 ml) to prepare a solution. A solution of triphosgene (70 mg) in chloroform (0.4 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1R)-1-(3-bromophenyl)ethylamine (70 mg) in chloroform (0.4 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (17 mg, yield 12%).

[0349] ¹H-NMR (CDCl₃): 1.47 (3H, d, J = 6.83 Hz), 2.17 (3H, s), 4.03 (3H, s), 4.04 (3H, s), 4.96 (1H, m), 5.19 (1H, d, J = 7.08 Hz), 6.25 (1H, s), 6.48 (1H, d, J = 5.12 Hz), 7.01 (1H, s), 7.03 (1H, overlapped), 7.21 (1H, dd, J = 7.81, 7.81 Hz), 7.27 (1H, overlapped), 7.38 (1H, d, J = 8.29 Hz), 7.42 (1H, s), 7.48 (1H, s), 7.52 (1H, s), 7.62 (1H, d, J = 9.51 Hz), 8.41 (1H, d, J = 5.12 Hz)

[0350] Mass spectrometric value (ESI-MS, m/z): 534, 536 (M-1)

[0351] $[\alpha]_D^{24} - 17.0^\circ$ (c0.5, CHCl₃)

Example 59: N-[(1R)-1-(3-Bromophenyl)ethyl]-N'-[4-[(6,7-dimethoxy-4-quinolyl)oxy]-2-methoxyphenyl]urea

[0352] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methoxyaniline (91 mg) was dissolved in chloroform (2 ml) and triethylamine (0.2 ml) to prepare a solution. A solution of triphosgene (70 mg) in chloroform (0.4 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1R)-1-(3-bromophenyl)ethylamine (70 mg) in chloroform (0.4 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (23 mg, yield 15%).

[0353] ¹H-NMR (CDCl₃): 1.51 (3H, d, J = 6.83 Hz), 3.76 (3H, s), 4.04 (3H, s), 4.05 (3H, s), 4.94 (1H, m), 5.14 (1H, d, J = 6.59 Hz), 6.45 (1H, d, J = 5.12 Hz), 6.67 (1H, d, J = 2.68 Hz), 6.77 (1H, dd, J = 8.78, 2.44 Hz), 6.87 (1H, s), 7.23 (1H, dd, J = 7.81, 7.81 Hz), 7.33 (1H, d, J = 7.81 Hz), 7.41 (1H, overlapped), 7.42 (1H, s), 7.53 (1H, s), 7.55 (1H, s),

8.16 (1H, d, J = 8.73 Hz), 8.46 (1H, d, J = 5.12 Hz)

[0354] Mass spectrometric value (ESI-MS, m/z): 550, 552 (M-1)

[0355] $[\alpha]_D^{24+16.2^\circ}$ (c0.5, CHCl₃)

5 Example 60: N-[(1R)-1-(3-Bromophenyl)ethyl]-N'-[4-[(6,7-dimethoxy-4-quinolyl)oxy]-2-(trifluoromethyl)phenyl]urea

[0356] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-(trifluoromethyl)aniline (82 mg) was dissolved in chloroform (2 ml) and triethylamine (0.2 ml) to prepare a solution. A solution of triphosgene (70 mg) in chloroform (0.4 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1R)-1-(3-bromophenyl)ethylamine (70 mg) in chloroform (0.4 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (39 mg, yield 29%).

[0357] ¹H-NMR (CDCl₃): 1.50 (3H, d, J = 7.08 Hz), 4.04 (3H, s), 4.05 (3H, s), 4.94 (1H, m), 5.51 (1H, d, J = 6.59 Hz), 6.45 (1H, d, J = 5.37 Hz), 6.72 (1H, s), 7.22 (1H, dd, J = 7.81, 7.56 Hz), 7.28 - 7.34 (2H, overlapped), 7.39 - 7.41 (2H, overlapped), 7.44 (1H, s), 7.50 (2H, s), 8.13 (1H, d, J = 9.03 Hz), 8.50 (1H, d, J = 5.37 Hz)

[0358] Mass spectrometric value (ESI-MS, m/z): 588, 590 (M-1)

[0359] $[\alpha]_D^{24+42.6^\circ}$ (c0.5, CHCl₃)

20 Example 61: N-[(1R)-1-(3-Bromophenyl)ethyl]-N'-[4-[(6,7-dimethoxy-4-quinolyl)oxy]-3-methoxyphenyl]urea

[0360] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-3-methoxyaniline (79 mg) was dissolved in chloroform (2 ml) and triethylamine (0.2 ml) to prepare a solution. A solution of triphosgene (70 mg) in chloroform (0.4 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1R)-1-(3-bromophenyl)ethylamine (70 mg) in chloroform (0.4 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (39 mg, yield 29%).

[0361] ¹H-NMR (CDCl₃): 1.46 (3H, d, J = 6.83 Hz), 3.70 (3H, s), 4.01 (3H, s), 4.04 (3H, s), 4.98 (1H, m), 5.76 (1H, d, J = 7.32 Hz), 6.30 (1H, d, J = 5.37 Hz), 6.67 (1H, dd, J = 8.54, 2.44 Hz), 6.99 (1H, d, J = 8.54 Hz), 7.17 (1H, dd, J = 7.81, 7.56 Hz), 7.27 (1H, overlapped), 7.35 (1H, d, J = 8.05 Hz), 7.40 (1H, s), 7.46 - 7.48 (3H, overlapped), 7.61 (1H, s), 8.39 (1H, d, J = 5.37 Hz)

[0362] Mass spectrometric value (ESI-MS, m/z): 550, 552 (M-1)

[0363] $[\alpha]_D^{24-28.0^\circ}$ (c0.5, CHCl₃)

35 Example 62: N-[(1R)-1-(3-Bromophenyl)ethyl]-N'-[4-[(6,7-dimethoxy-4-quinolyl)oxy]-2,3-dimethylphenyl]urea

[0364] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2,3-dimethylaniline (79 mg) was dissolved in chloroform (2 ml) and triethylamine (0.2 ml) to prepare a solution. A solution of triphosgene (70 mg) in chloroform (0.4 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1R)-1-(3-bromophenyl)ethylamine (70 mg) in chloroform (0.4 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (34 mg, yield 26%).

[0365] ¹H-NMR (CDCl₃): 1.46 (3H, d, J = 7.07 Hz), 2.11 (3H, s), 2.18 (3H, s), 4.05 (3H, s), 4.06 (3H, s), 4.99 (1H, m), 5.12 (1H, d, J = 7.32 Hz), 6.27 (1H, d, J = 5.37 Hz), 6.31 (1H, s), 6.99 (1H, d, J = 8.72 Hz), 7.20 (1H, dd, J = 7.81, 7.81 Hz), 7.27 (1H, overlapped), 7.35 - 7.39 (2H, overlapped), 7.45 (1H, s), 7.46 (1H, s), 7.59 (1H, s), 8.43 (1H, d, J = 5.37 Hz)

[0366] Mass spectrometric value (ESI-MS, m/z): 548, 550 (M-1)

[0367] $[\alpha]_D^{24-31.3^\circ}$ (c0.5, CHCl₃)

50 Example 63: N-[(1R)-1-(3-Bromophenyl)ethyl]-N'-[4-[(6,7-dimethoxy-4-quinolyl)oxy]-2-fluorophenyl]urea

[0368] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-fluoroaniline (79 mg) was dissolved in chloroform (2 ml) and triethylamine (0.2 ml) to prepare a solution. A solution of triphosgene (70 mg) in chloroform (0.4 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1R)-1-(3-bromophenyl)ethylamine (70 mg) in chloroform (0.4 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (58 mg, yield 43%).

[0369] ¹H-NMR (CDCl₃): 1.48 (3H, d, J = 7.07 Hz), 4.02 (3H, s), 4.03 (3H, s), 4.96 (1H, m), 5.76 (1H, d, J = 6.83 Hz), 6.47 (1H, d, J = 5.37 Hz), 6.89 (1H, dd, J = 11.47, 2.44 Hz), 6.94 (1H, dd, J = 8.78, 2.44 Hz), 7.05 (1H, brs), 7.19

(1H, dd, J = 7.81, 7.81 Hz), 7.28 (1H, d, J = 7.81 Hz), 7.37 (1H, ddd, J = 7.81, 1.71, 1.22 Hz), 7.42 (1H, s), 7.49 (1H, dd, J = 1.71, 1.71 Hz), 7.50 (1H, s), 8.18 (1H, dd, J = 9.03, 9.03 Hz), 8.47 (1H, d, J = 5.37 Hz)

[0370] Mass spectrometric value (ESI-MS, m/z): 538, 540 (M-1)

[0371] $[\alpha]_D^{24} + 53.4^\circ$ (c0.5, CHCl₃)

Example 64: N-(3-Fluoro-4-([6-methoxy-7-(3-morpholinopropoxy)-4-quinolyl]oxy)phenyl)-N'-[1-(4-fluorophenyl)ethyl]urea

[0372] 3-Fluoro-4-([6-methoxy-7-(3-morpholinopropoxy)-4-quinolyl]oxy)-aniline (30 mg) was dissolved in chloroform (3 ml) and triethylamine (0.3 ml) to prepare a solution. A solution of triphosgene (17 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for 25 min. Next, a solution of 1-(4-fluorophenyl)ethylamine (12 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 5 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (22 mg, yield 52%).

[0373] ¹H-NMR (CDCl₃, 400 MHz): δ 1.50 (d, J = 6.8 Hz, 3H), 2.09 - 2.15 (m, 2H), 2.48 (br, 4H), 2.57 (t, J = 7.1 Hz, 2H), 3.71 - 3.73 (m, 4H), 4.02 (s, 3H), 4.24 (t, J = 6.6 Hz, 2H), 4.96 - 5.00 (m, 1H), 5.28 (br, 1H), 6.37 (d, J = 4.4 Hz, 1H), 7.00 - 7.05 (m, 3H), 7.12 (d, J = 8.5 Hz, 1H), 7.32 (dd, J = 5.4, 8.5 Hz, 1H), 7.41 (s, 1H), 7.48 (dd, J = 2.4, 12.2 Hz, 1H), 7.56 (s, 1H), 8.44 (d, J = 5.1 Hz, 1H)

[0374] Mass spectrometric value (ESI-MS, m/z): 593 (M⁺+1)

Example 65: N-[1-(3,4-Difluorophenyl)ethyl]-N'-(3-fluoro-4-([6-methoxy-7-(3-morpholinopropoxy)-4-quinolyl]oxy)phenyl)urea

[0375] 3-Fluoro-4-([6-methoxy-7-(3-morpholinopropoxy)-4-quinolyl]oxy)-aniline (30 mg) was dissolved in chloroform (3 ml) and triethylamine (0.3 ml) to prepare a solution. A solution of triphosgene (17 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for 25 min. Next, a solution of 1-(3,4-difluorophenyl)ethylamine (14 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 5 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (12 mg, yield 29%).

[0376] ¹H-NMR (CDCl₃, 400 MHz): δ 1.47 (d, J = 6.8 Hz, 3H), 2.08 - 2.15 (m, 2H), 2.48 (br, 4H), 2.57 (t, J = 7.1 Hz, 2H), 3.72 (t, J = 4.4 Hz, 4H), 4.03 (s, 3H), 4.24 (t, J = 6.6 Hz, 2H), 4.95 - 5.00 (m, 1H), 5.34 (br, 1H), 6.37 (d, J = 4.4 Hz, 1H), 7.02 - 7.22 (m, 5H), 7.41 (s, 1H), 7.49 (dd, J = 2.4, 12.2 Hz, 1H), 7.57 (s, 1H), 8.45 (d, J = 5.4 Hz, 1H)

[0377] Mass spectrometric value (ESI-MS, m/z): 609 (M⁺-1)

Example 66: N-[1-(2,4-Difluorophenyl)ethyl]-N'-(3-fluoro-4-([6-methoxy-7-(3-morpholinopropoxy)-4-quinolyl]oxy)phenyl)urea

[0378] 3-Fluoro-4-([6-methoxy-7-(3-morpholinopropoxy)-4-quinolyl]oxy)-aniline (30 mg) was dissolved in chloroform (3 ml) and triethylamine (0.3 ml) to prepare a solution. A solution of triphosgene (17 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for 25 min. Next, a solution of 1-(2,4-difluorophenyl)ethylamine (14 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 5 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (25 mg, yield 57%).

[0379] ¹H-NMR (CDCl₃, 400 MHz): δ 1.54 (d, J = 6.6 Hz, 3H), 2.11 - 2.16 (m, 2H), 2.49 (br, 4H), 2.58 (t, J = 6.8 Hz, 2H), 3.73 (t, J = 4.6 Hz, 4H), 4.03 (s, 3H), 4.26 (t, J = 6.8 Hz, 2H), 5.13 - 5.19 (m, 1H), 6.37 (d, J = 5.4 Hz, 1H), 6.57 (s, 1H), 6.79 - 6.88 (m, 2H), 6.97 - 7.06 (m, 1H), 7.14 (t, J = 8.8 Hz, 1H), 7.29 - 7.35 (m, 1H), 7.42 (s, 1H), 7.49 (dd, J = 2.4, 12.2 Hz, 1H), 7.56 (s, 1H), 8.45 (d, J = 5.4 Hz, 1H)

[0380] Mass spectrometric value (ESI-MS, m/z): 609 (M⁺-1)

Example 67: N-(2-Fluoro-4-([6-methoxy-7-(3-morpholinopropoxy)-4-quinolyl]oxy)phenyl)-N'-[1-(4-fluorophenyl)ethyl]urea

[0381] 2-Fluoro-4-([6-methoxy-7-(3-morpholinopropoxy)-4-quinolyl]oxy)-aniline (30 mg) was dissolved in chloroform (3 ml) and triethylamine (0.3 ml) to prepare a solution. A solution of triphosgene (17 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for 25 min. Next, a solution of 1-(4-fluorophenyl)ethylamine (12 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 5 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (31 mg, yield 73%).

[0382] ¹H-NMR (CDCl₃, 400 MHz): δ 1.53 (d, J = 6.8 Hz, 3H), 2.09 - 2.16 (m, 2H), 2.48 (br, 4H), 2.57 (t, J = 7.1 Hz, 2H), 3.72 (t, J = 4.9 Hz, 4H), 4.01 (s, 3H), 4.26 (t, J = 6.8 Hz, 2H), 4.95 - 4.99 (m, 1H), 5.04 (d, J = 6.8 Hz, 1H), 6.46 (d, J = 5.1 Hz, 1H), 6.48 (d, J = 2.9 Hz, 1H), 6.89 - 6.96 (m, 2H), 7.05 (t, J = 8.5 Hz, 2H), 7.35 (dd, J = 5.4, 8.8 Hz, 2H), 7.42 (s, 1H), 7.48 (s, 1H), 8.16 (t, J = 9.0 Hz, 1H), 8.48 (d, J = 5.4 Hz, 1H)

[0383] Mass spectrometric value (ESI-MS, m/z): 591 (M⁺-1)

Example 68: N-[1-(3,4-Difluorophenyl)ethyl]-N'-(2-fluoro-4-([6-methoxy-7-(3-morpholinopropoxy)-4-quinolyl]oxy)phenyl)urea

[0384] 2-Fluoro-4-([6-methoxy-7-(3-morpholinopropoxy)-4-quinolyl]oxy)-aniline (30 mg) was dissolved in chloroform (3 ml) and triethylamine (0.3 ml) to prepare a solution. A solution of triphosgene (17 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for 25 min. Next, a solution of 1-(3,4-difluorophenyl)ethylamine (14 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 5 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (27 mg, yield 60%).

[0385] ¹H-NMR (CDCl₃, 400 MHz): δ 1.52 (d, J = 6.8 Hz, 3H), 2.09 - 2.16 (m, 2H), 2.49 (br, 4H), 2.58 (t, J = 7.1 Hz, 2H), 3.73 (t, J = 4.6 Hz, 4H), 4.01 (s, 3H), 4.26 (t, J = 6.6 Hz, 2H), 4.95 - 5.00 (m, 1H), 5.09 (d, J = 6.8 Hz, 1H), 6.47 (d, J = 5.4 Hz, 1H), 6.54 (d, J = 3.2 Hz, 1H), 6.91 - 6.96 (m, 2H), 7.09 - 7.23 (m, 3H), 7.42 (s, 1H), 7.48 (s, 1H), 8.14 (t, J = 8.8 Hz, 2H), 8.48 (d, J = 5.1 Hz, 1H)

[0386] Mass spectrometric value (ESI-MS, m/z): 609 (M⁺-1)

Example 69: N-[1-(2,4-Difluorophenyl)ethyl]-N'-(2-fluoro-4-([6-methoxy-7-(3-morpholinopropoxy)-4-quinolyl]oxy)phenyl)urea

[0387] 2-Fluoro-4-([6-methoxy-7-(3-morpholinopropoxy)-4-quinolyl]oxy)-aniline (30 mg) was dissolved in chloroform (3 ml) and triethylamine (0.3 ml) to prepare a solution. A solution of triphosgene (17 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for 25 min. Next, a solution of 1-(2,4-difluorophenyl)ethylamine (14 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 5 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (29 mg, yield 66%).

[0388] ¹H-NMR (CDCl₃, 400 MHz): δ 1.55 (d, J = 6.6 Hz, 3H), 2.09 - 2.16 (m, 2H), 2.49 (br, 4H), 2.58 (t, J = 6.8 Hz, 2H), 3.73 (t, J = 4.6 Hz, 4H), 4.01 (s, 3H), 4.27 (t, J = 6.8 Hz, 2H), 5.14 - 5.17 (m, 1H), 5.22 (d, J = 7.8 Hz, 1H), 6.46 (d, J = 5.1 Hz, 1H), 6.50 (d, J = 2.7 Hz, 1H), 6.80 - 6.88 (m, 2H), 6.91 - 7.00 (m, 2H), 7.33 (dd, J = 8.5, 14.6 Hz, 1H), 7.43 (s, 1H), 7.48 (s, 1H), 8.14 (t, J = 8.8 Hz, 2H), 8.48 (d, J = 5.4 Hz, 1H)

[0389] Mass spectrometric value (ESI-MS, m/z): 609 (M⁺-1)

Example 70: N-[4-([6,7-Dimethoxy-4-quinolyl]oxy)-2-fluorophenyl]-N'-[1-(1,3-thiazol-2-yl)ethyl]urea

[0390] 4-([6,7-Dimethoxy-4-quinolyl]oxy)-2-fluoroaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (35 mg, yield 79%).

[0391] ¹H-NMR (CDCl₃, 400 MHz): δ 1.70 (d, J = 7.1 Hz, 3H), 4.04 (s, 3H), 4.05 (s, 3H), 5.37 - 5.42 (m, 1H), 5.80 (d, J = 7.3 Hz, 1H), 6.49 (d, J = 5.1 Hz, 1H), 6.93 - 6.98 (m, 2H), 7.01 (br, 1H), 7.31 (d, J = 3.0 Hz, 1H), 7.42 (s, 1H), 7.50 (s, 1H), 7.74 (d, J = 3.2 Hz, 1H), 8.17 (d, J = 9.3 Hz, 1H), 8.50 (d, J = 5.4 Hz, 1H)

[0392] Mass spectrometric value (ESI-MS, m/z): 469 (M⁺+1)

Example 71: N-[4-([6,7-Dimethoxy-4-quinolyl]oxy)-2-fluorophenyl]-N'-[(1S)-1-(1,3-thiazol-2-yl)ethyl]urea

[0393] 4-([6,7-Dimethoxy-4-quinolyl]oxy)-2-fluoroaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1S)-1-(1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (30 mg, yield 66%).

[0394] ¹H-NMR (CDCl₃, 400 MHz): δ 1.71 (d, J = 6.8 Hz, 3H), 4.07 (s, 3H), 4.10 (s, 3H), 5.34 - 5.39 (m, 1H), 6.60

(d, J = 6.1 Hz, 1H), 6.61 (s, 1H), 6.90 (dd, J = 2.7, 11.0 Hz, 1H), 6.97 - 6.99 (m, 1H), 7.27 (s, 1H), 7.57 (s, 1H), 7.70 (d, J = 3.4 Hz, 1H), 7.74 (br, 1H), 7.77 (s, 1H), 8.33 (dd, J = 9.0, 9.0 Hz, 1H), 8.47 (d, J = 6.1 Hz, 1H)

[0395] Mass spectrometric value (ESI-MS, m/z): 469 (M⁺+1)

[0396] $[\alpha]_D^{25} -39.2^\circ$ (c0.5, CHCl₃)

Example 72: N-{4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-fluorophenyl}-N'-(1R)-1-(1,3-thiazol-2-yl)ethylurea

[0397] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-fluoroaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1R)-1-(1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (28 mg, yield 62%).

[0398] ¹H-NMR (CDCl₃, 400 MHz): δ 1.70 (d, J = 7.1 Hz, 3H), 4.06 (s, 3H), 4.07 (s, 3H), 5.35 - 5.42 (m, 1H), 6.48 (d, J = 6.8 Hz, 1H), 6.55 (d, J = 5.9 Hz, 1H), 6.91 (dd, J = 2.7, 11.2 Hz, 1H), 6.94 - 6.98 (m, 1H), 7.28 (d, J = 3.2 Hz, 1H), 7.54 (s, 1H), 7.60 (br, 1H), 7.63 (s, 1H), 7.71 (d, J = 3.4 Hz, 1H), 8.26 (d, J = 9.0 Hz, 1H), 8.48 (d, J = 5.6 Hz, 1H)

[0399] Mass spectrometric value (ESI-MS, m/z): 469 (M⁺+1)

[0400] $[\alpha]_D^{25} +40.7^\circ$ (c0.5, CHCl₃)

Example 73: N-{4-[(6,7-Dimethoxy-4-quinolyl)oxy]-3-fluorophenyl}-N'-(1-(1,3-thiazol-2-yl)ethyl)urea

[0401] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-3-fluoroaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (39 mg, yield 87%).

[0402] ¹H-NMR (CDCl₃, 400 MHz): δ 1.69 (d, J = 6.8 Hz, 3H), 4.03 (s, 3H), 4.04 (s, 3H), 5.33 - 5.42 (m, 1H), 5.95 (d, J = 7.3 Hz, 1H), 6.39 (dd, J = 1.0, 5.1 Hz, 1H), 7.05 (dd, J = 1.5, 8.8 Hz, 1H), 7.13 (d, J = 8.8, 8.8 Hz, 1H), 7.34 (d, J = 3.2 Hz, 1H), 7.41 (s, 1H), 7.53 (d, J = 2.4, 12.2 Hz, 1H), 7.57 (s, 1H), 7.74 (d, J = 3.2 Hz, 1H), 8.47 (d, J = 5.4 Hz, 1H)

[0403] Mass spectrometric value (ESI-MS, m/z): 469 (M⁺+1)

Example 74: N-{4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methoxyphenyl}-N'-(1-(1,3-thiazol-2-yl)ethyl)urea

[0404] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methoxyaniline (816 mg) was dissolved in chloroform (40 ml) and triethylamine (2.0 ml) to prepare a solution. A solution of triphosgene (368 mg) in chloroform (1.0 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(1,3-thiazol-2-yl)-1-ethylamine (352 mg) in chloroform (1.0 ml) was added thereto, and the mixture was stirred at room temperature overnight. Water (40 ml) was added thereto, and the mixture was stirred for 10 min, followed by extraction with chloroform. The organic layer was washed with saturated brine and was dried over sodium sulfate. The solvent was removed by evaporation, and the residue was purified by chromatography on silica gel using chloroform/acetone for development to give the title compound (860 mg, 72%).

[0405] ¹H-NMR (CDCl₃, 400 MHz): δ 1.69 (d, J = 6.8 Hz, 3H), 3.83 (s, 3H), 4.06 (s, 3H), 4.07 (s, 3H), 5.37 - 5.45 (m, 1H), 5.67 (d, J = 7.3 Hz, 1H), 6.51 (d, J = 5.4 Hz, 1H), 6.70 (d, J = 2.4 Hz, 1H), 6.79 (dd, J = 2.4, 8.8 Hz, 1H), 7.02 (s, 1H), 7.30 (d, J = 3.4 Hz, 1H), 7.53 (s, 1H), 7.57 (s, 1H), 7.73 (d, J = 3.4 Hz, 1H), 8.19 (d, J = 8.8 Hz, 1H), 8.47 (d, J = 5.4 Hz, 1H)

[0406] Mass spectrometric value (ESI-MS, m/z): 481 (M⁺+1)

Example 75: N-{4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methoxyphenyl}-N'-(1S)-1-(1,3-thiazol-2-yl)ethyl)urea

[0407] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methoxyaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1S)-1-(1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (33 mg, yield 75%).

[0408] ¹H-NMR (CDCl₃, 400 MHz): δ 1.68 (d, J = 6.8 Hz, 3H), 3.78 (s, 3H), 4.05 (s, 3H), 4.07 (s, 3H), 5.38 - 5.45 (m, 1H), 6.08 (d, J = 7.6 Hz, 1H), 6.52 (d, J = 5.6 Hz, 1H), 6.67 (d, J = 2.4 Hz, 1H), 6.77 (dd, J = 2.7, 8.8 Hz, 1H), 7.21 (s,

1H), 7.30 (d, J = 3.4 Hz, 1H), 7.57 (s, 1H), 7.59 (s, 1H), 7.72 (d, J = 3.2 Hz, 1H), 8.21 (d, J = 8.8 Hz, 1H), 8.46 (d, J = 5.6 Hz, 1H)

[0409] Mass spectrometric value (ESI-MS, m/z): 481 (M⁺+1)

[0410] $[\alpha]_D^{25}$ -16.4° (c0.5, CHCl₃)

Example 76: N-{4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methoxyphenyl}-N'-[(1R)-1-(1,3-thiazol-2-yl)ethyl]urea

[0411] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methoxyaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1R)-1-(1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (36 mg, yield 82%).

[0412] ¹H-NMR (CDCl₃, 400 MHz): δ 1.68 (d, J = 6.8 Hz, 3H), 3.78 (s, 3H), 4.05 (s, 3H), 4.07 (s, 3H), 5.38 - 5.45 (m, 1H), 6.08 (d, J = 7.6 Hz, 1H), 6.52 (d, J = 5.6 Hz, 1H), 6.67 (d, J = 2.4 Hz, 1H), 6.77 (dd, J = 2.7, 8.8 Hz, 1H), 7.21 (s, 1H), 7.30 (d, J = 3.4 Hz, 1H), 7.57 (s, 1H), 7.59 (s, 1H), 7.72 (d, J = 3.2 Hz, 1H), 8.21 (d, J = 8.8 Hz, 1H), 8.46 (d, J = 5.6 Hz, 1H)

[0413] Mass spectrometric value (ESI-MS, m/z): 481 (M⁺+1)

[0414] $[\alpha]_D^{24}$ +14.1° (c0.5, CHCl₃)

Example 77: N-{4-[(6,7-Dimethoxy-4-quinolyl)oxy]phenyl}-N'-[(1,3-thiazol-2-yl)ethyl]urea

[0415] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]aniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (36 mg, yield 78%).

[0416] ¹H-NMR (CDCl₃, 400 MHz): δ 1.69 (d, J = 7.1 Hz, 3H), 4.06 (s, 3H), 4.07 (s, 3H), 5.34 - 5.40 (m, 1H), 6.51 (d, J = 6.6 Hz, 1H), 6.70 (s, 1H), 7.05 (d, J = 9.0 Hz, 1H), 7.20 (d, J = 3.2 Hz, 1H), 7.64 (s, 1H), 7.65 (d, J = 3.2 Hz, 1H), 7.72 (d, J = 9.0 Hz, 1H), 7.89 (s, 1H), 8.26 (d, J = 6.6 Hz, 1H), 8.91 (s, 1H)

[0417] Mass spectrometric value (ESI-MS, m/z): 451 (M⁺+1)

Example 78: N-{4-[(6,7-Dimethoxy-4-quinolyl)oxy]phenyl}-N'-[(1S)-1-(1,3-thiazol-2-yl)ethyl]urea

[0418] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]aniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1S)-1-(1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (31 mg, yield 68%).

[0419] ¹H-NMR (CDCl₃, 400 MHz): δ 1.68 (d, J = 6.8 Hz, 3H), 4.09 (s, 3H), 4.10 (s, 3H), 5.32 - 5.38 (m, 1H), 6.51 (d, J = 6.3 Hz, 1H), 6.55 (br, 1H), 7.06 (d, J = 8.8 Hz, 1H), 7.22 (d, J = 3.2 Hz, 1H), 7.64 (d, J = 1.2 Hz, 1H), 7.66 (d, J = 3.4 Hz, 1H), 7.67 (s, 1H), 7.79 (s, 1H), 8.31 (d, J = 6.3 Hz, 1H), 8.59 (s, 1H)

[0420] Mass spectrometric value (ESI-MS, m/z): 451 (M⁺+1)

[0421] $[\alpha]_D^{25}$ -24.6° (c0.5, CHCl₃)

Example 79: N-{4-[(6,7-Dimethoxy-4-quinolyl)oxy]phenyl}-N'-[(1R)-1-(1,3-thiazol-2-yl)ethyl]urea

[0422] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]aniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1R)-1-(1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (28 mg, yield 62%).

[0423] ¹H-NMR (CDCl₃, 400 MHz): δ 1.69 (d, J = 7.1 Hz, 3H), 4.11 (s, 3H), 4.12 (s, 3H), 5.31 - 5.37 (m, 1H), 6.51 (d, J = 6.6 Hz, 1H), 6.70 (s, 1H), 7.05 (d, J = 9.0 Hz, 1H), 7.20 (d, J = 3.2 Hz, 1H), 7.65 (s, 1H), 7.65 (d, J = 3.2 Hz, 1H), 7.72 (d, J = 9.0 Hz, 1H), 7.89 (s, 1H), 8.26 (d, J = 6.6 Hz, 1H), 8.91 (s, 1H)

[0424] Mass spectrometric value (ESI-MS, m/z): 451 (M^{+1})

[0425] $[\alpha]_D^{24+23.9^\circ}$ (c0.5, CHCl_3)

Example 80: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-3-methoxyphenyl]-N'-[1-(1,3-thiazol-2-yl)ethyl]urea

[0426] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-3-methoxyaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (28 mg, yield 62%).

[0427] $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 1.69 (d, J = 6.8 Hz, 3H), 3.71 (s, 3H), 4.09 (s, 3H), 4.10 (s, 3H), 5.34 - 5.39 (m, 1H), 6.38 (d, J = 6.3 Hz, 1H), 6.70 (br, 1H), 6.94 (dd, J = 2.2, 8.5 Hz, 1H), 6.99 (d, J = 8.8 Hz, 1H), 7.20 (d, J = 3.2 Hz, 1H), 7.66 (d, J = 2.9 Hz, 1H), 7.66 (s, 1H), 7.73 (d, J = 2.2 Hz, 1H), 7.80 (s, 1H), 8.23 (d, J = 6.6 Hz, 1H), 8.78 (br, 1H)

[0428] Mass spectrometric value (ESI-MS, m/z): 479 (M^{+1})

Example 81: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-3-methoxyphenyl]-N'-[(1S)-1-(1,3-thiazol-2-yl)ethyl]urea

[0429] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-3-methoxyaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1S)-1-(1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (31 mg, yield 72%).

[0430] $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 1.70 (d, J = 6.8 Hz, 3H), 3.70 (s, 3H), 4.11 (s, 3H), 4.13 (s, 3H), 5.33 - 5.39 (m, 1H), 6.39 (d, J = 6.6 Hz, 1H), 6.87 (s, 1H), 6.99 (s, 1H), 7.18 (d, J = 3.2 Hz, 1H), 7.65 (d, J = 3.4 Hz, 1H), 7.68 (s, 1H), 7.82 (s, 1H), 7.92 (s, 1H), 8.17 (d, J = 6.6 Hz, 1H), 9.16 (s, 1H)

[0431] Mass spectrometric value (ESI-MS, m/z): 479 (M^{+1})

[0432] $[\alpha]_D^{25+24.0^\circ}$ (c0.5, CHCl_3)

Example 82: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-3-methoxyphenyl]-N'-[(1R)-1-(1,3-thiazol-2-yl)ethyl]urea

[0433] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-3-methoxyaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1R)-1-(1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (31 mg, yield 71%).

[0434] $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 1.68 (d, J = 6.8 Hz, 3H), 3.71 (s, 3H), 4.06 (s, 3H), 4.07 (s, 3H), 5.35 - 5.43 (m, 1H), 6.36 (d, J = 6.1 Hz, 1H), 6.48 (s, 1H), 6.86 (dd, J = 2.2, 8.5 Hz, 1H), 7.00 (d, J = 8.5 Hz, 1H), 7.25 (d, J = 3.4 Hz, 1H), 7.59 (d, J = 2.2 Hz, 1H), 7.62 (s, 1H), 7.63 (s, 1H), 7.67 (d, J = 3.2 Hz, 1H), 8.26 (s, 1H), 8.33 (d, J = 5.9 Hz, 1H)

[0435] Mass spectrometric value (ESI-MS, m/z): 479 (M^{+1})

[0436] $[\alpha]_D^{25+26.7^\circ}$ (c0.5, CHCl_3)

Example 83: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-(trifluoromethyl)phenyl]-N'-[1-(1,3-thiazol-2-yl)ethyl]urea

[0437] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-(trifluoromethyl)aniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (29 mg, yield 68%).

[0438] $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 1.71 (d, J = 7.1 Hz, 3H), 4.08 (s, 3H), 4.12 (s, 3H), 5.34 - 5.40 (m, 1H), 6.29 (br, 1H), 6.61 (d, J = 6.1 Hz, 1H), 7.21 (s, 1H), 7.30 (d, J = 3.4 Hz, 1H), 7.38 (dd, J = 2.9, 9.0 Hz, 1H), 7.44 (d, J = 2.9 Hz, 1H), 7.57 (s, 1H), 7.72 (d, J = 3.2 Hz, 1H), 7.87 (br, 1H), 8.27 (d, J = 9.0 Hz, 1H), 8.50 (d, J = 6.1 Hz, 1H)

[0439] Mass spectrometric value (ESI-MS, m/z): 517 (M^{+1})

Example 84: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-(trifluoromethyl)phenyl]-N'-[(1S)-1-(1,3-thiazol-2-yl)ethyl]urea

[0440] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-(trifluoromethyl)aniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1S)-1-(1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (29 mg, yield 68%).

[0441] ¹H-NMR (CDCl₃, 400 MHz): δ 1.70 (d, J = 6.8 Hz, 3H), 4.07 (s, 3H), 4.10 (s, 3H), 5.34 - 5.41 (m, 1H), 6.26 (d, J = 6.8 Hz, 1H), 6.58 (d, J = 5.9 Hz, 1H), 7.16 (s, 1H), 7.30 (d, J = 3.2 Hz, 1H), 7.37 (dd, J = 2.9, 9.0 Hz, 1H), 7.44 (d, J = 2.7 Hz, 1H), 7.55 (s, 1H), 7.72 (d, J = 3.4 Hz, 1H), 7.75 (br, 1H), 8.23 (d, J = 9.0 Hz, 1H), 8.51 (d, J = 5.9 Hz, 1H)

[0442] Mass spectrometric value (ESI-MS, m/z): 517 (M⁺-1)

[0443] [α]_D²⁵-34.3° (c0.5, CHCl₃)

Example 85: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-(trifluoromethyl)phenyl]-N'-[(1R)-1-(1,3-thiazol-2-yl)ethyl]urea

[0444] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-(trifluoromethyl)aniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1R)-1-(1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (30 mg, yield 71%).

[0445] ¹H-NMR (CDCl₃, 400 MHz): δ 1.70 (d, J = 6.8 Hz, 3H), 4.06 (s, 3H), 4.09 (s, 3H), 5.35 - 5.42 (m, 1H), 6.33 (d, J = 7.1 Hz, 1H), 6.55 (d, J = 5.9 Hz, 1H), 7.17 (s, 1H), 7.30 (d, J = 3.2 Hz, 1H), 7.36 (dd, J = 2.9, 9.0 Hz, 1H), 7.43 (d, J = 2.9 Hz, 1H), 7.54 (s, 1H), 7.67 (s, 1H), 7.71 (d, J = 3.4 Hz, 1H), 8.20 (d, J = 9.0 Hz, 1H), 8.51 (d, J = 5.9 Hz, 1H)

[0446] Mass spectrometric value (ESI-MS, m/z): 517 (M⁺-1)

[0447] [α]_D²⁵+36.6° (c0.5, CHCl₃)

Example 86: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2,5-dimethylphenyl]-N'-[(1S)-1-(1,3-thiazol-2-yl)ethyl]urea

[0448] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2,5-dimethylaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1S)-1-(1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (30 mg, yield 68%).

[0449] ¹H-NMR (CDCl₃, 400 MHz): δ 1.68 (d, J = 6.8 Hz, 3H), 2.10 (s, 3H), 2.30 (s, 3H), 4.09 (s, 3H), 4.11 (s, 3H), 5.33 - 5.39 (m, 1H), 6.29 (br, 1H), 6.47 (d, J = 6.1 Hz, 1H), 6.92 (s, 1H), 7.16 (br, 1H), 7.63 (s, 1H), 7.67 (d, J = 3.2 Hz, 1H), 7.75 (s, 1H), 7.84 (s, 1H), 8.38 (d, J = 6.1 Hz, 1H)

[0450] Mass spectrometric value (ESI-MS, m/z): 479 (M⁺+1)

[0451] [α]_D²⁵+10.6° (c0.5, CHCl₃)

Example 87: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2,5-dimethylphenyl]-N'-[(1R)-1-(1,3-thiazol-2-yl)ethyl]urea

[0452] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2,5-dimethylaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of (1R)-1-(1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (30 mg, yield 68%).

[0453] ¹H-NMR (CDCl₃, 400 MHz): δ 1.67 (d, J = 6.8 Hz, 3H), 2.12 (s, 3H), 2.25 (s, 3H), 4.07 (s, 3H), 4.07 (s, 3H), 5.35 - 5.42 (m, 1H), 6.13 (d, J = 7.1 Hz, 1H), 6.38 (d, J = 5.6 Hz, 1H), 6.92 (s, 1H), 6.98 (s, 1H), 7.27 (d, J = 3.2 Hz, 1H), 7.60 (s, 1H), 7.65 (s, 1H), 7.68 (d, J = 3.4 Hz, 1H), 8.42 (d, J = 5.6 Hz, 1H)

[0454] Mass spectrometric value (ESI-MS, m/z): 479 (M⁺+1)

[0455] [α]_D²⁵-9.0° (c0.5, CHCl₃)

Example 88: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-fluorophenyl]-N'-[1-(4-methyl-1,3-thiazol-2-yl)ethyl]urea

[0456] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-fluoroaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(4-methyl-1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (38 mg, yield 83%).

[0457] ¹H-NMR (CDCl₃, 400 MHz): δ 1.68 (d, J = 7.1 Hz, 3H), 2.43 (d, J = 1.0 Hz, 3H), 4.06 (s, 3H), 4.08 (s, 3H), 5.26 - 5.33 (m, 1H), 6.38 (br, 1H), 6.57 (d, J = 5.9 Hz, 1H), 6.81 (d, J = 1.0 Hz, 1H), 6.91 (dd, J = 2.7, 11.0 Hz, 1H), 6.95 - 6.98 (m, 1H), 7.55 (s, 1H), 7.69 (s, 1H), 7.69 (br, 1H), 8.28 (t, J = 9.0 Hz, 1H), 8.48 (d, J = 5.9 Hz, 1H)

[0458] Mass spectrometric value (ESI-MS, m/z): 483 (M⁺+1)

Example 89: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-3-fluorophenyl]-N'-[1-(4-methyl-1,3-thiazol-2-yl)ethyl]urea

[0459] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-3-fluoroaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(4-methyl-1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (29 mg, yield 63%).

[0460] ¹H-NMR (CDCl₃, 400 MHz): δ 1.65 (d, J = 6.8 Hz, 3H), 2.40 (d, J = 0.7 Hz, 3H), 4.04 (s, 3H), 4.06 (s, 3H), 5.25 - 5.32 (m, 1H), 6.43 (br, 1H), 6.46 (d, J = 4.9 Hz, 1H), 6.80 (d, J = 1.2 Hz, 1H), 7.08 (t, J = 8.8 Hz, 1H), 7.14 (dd, J = 2.2, 8.8 Hz, 1H), 7.57 (s, 1H), 7.58 - 7.62 (m, 2H), 8.41 (d, J = 5.9 Hz, 1H), 8.44 (br, 1H)

[0461] Mass spectrometric value (ESI-MS, m/z): 483 (M⁺+1)

Example 90: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methoxyphenyl]-N'-[1-(4-methyl-1,3-thiazol-2-yl)ethyl]urea

[0462] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methoxyaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(4-methyl-1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (22 mg, yield 48%).

[0463] ¹H-NMR (CDCl₃, 400 MHz): δ 1.66 (d, J = 7.1 Hz, 3H), 2.44 (d, J = 1.0 Hz, 3H), 3.82 (s, 3H), 4.07 (s, 3H), 4.11 (s, 3H), 5.29 - 5.36 (m, 1H), 6.01 (d, J = 7.1 Hz, 1H), 6.60 (d, J = 6.1 Hz, 1H), 6.68 (d, J = 2.7 Hz, 1H), 6.78 (dd, J = 2.7, 9.0 Hz, 1H), 6.82 (d, J = 1.2 Hz, 1H), 7.23 (s, 1H), 7.60 (s, 1H), 7.84 (br, 1H), 8.28 (d, J = 8.8 Hz, 1H), 8.45 (d, J = 6.1 Hz, 1H)

[0464] Mass spectrometric value (ESI-MS, m/z): 495 (M⁺+1)

Example 91: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]phenyl]-N'-[1-(4-methyl-1,3-thiazol-2-yl)ethyl]urea

[0465] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methoxyaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(4-methyl-1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (32 mg, yield 68%).

[0466] ¹H-NMR (CDCl₃, 400 MHz): δ 1.65 (d, J = 6.8 Hz, 3H), 2.40 (d, J = 1.0 Hz, 3H), 4.06 (s, 3H), 4.07 (s, 3H), 5.25 - 5.32 (m, 1H), 6.24 (br, 1H), 6.48 (d, J = 5.6 Hz, 1H), 6.78 (d, J = 1.0 Hz, 1H), 7.06 (d, J = 9.0 Hz, 1H), 7.54 (d, J = 9.0 Hz, 1H), 7.58 (s, 1H), 7.60 (s, 1H), 8.09 (br, 1H), 8.39 (d, J = 5.9 Hz, 1H)

[0467] Mass spectrometric value (ESI-MS, m/z): 465 (M⁺+1)

Example 92: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-(trifluoromethyl)phenyl]-N'-[1-(4-methyl-1,3-thiazol-2-yl)ethyl]urea

[0468] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-(trifluoromethyl)aniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added

to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(4-methyl-1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (35 mg, yield 80%).

[0469] $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 1.67 (d, $J = 6.8$ Hz, 3H), 2.42 (d, $J = 1.0$ Hz, 3H), 4.06 (s, 3H), 4.08 (s, 3H), 5.27 - 5.34 (m, 1H), 6.33 (d, $J = 6.8$ Hz, 1H), 6.53 (d, $J = 5.6$ Hz, 1H), 6.82 (d, $J = 1.0$ Hz, 1H), 7.10 (br, 1H), 7.30 (dd, $J = 2.7, 9.0$ Hz, 1H), 7.42 (d, $J = 2.7$ Hz, 1H), 7.53 (s, 1H), 7.62 (s, 1H), 7.21 (d, $J = 9.0$ Hz, 1H), 8.51 (d, $J = 5.6$ Hz, 1H)
[0470] Mass spectrometric value (ESI-MS, m/z): 533 ($M^+ + 1$)

Example 93: N-{4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2,5-dimethylphenyl}-N'-[1-(4-methyl-1,3-thiazol-2-yl)ethyl]urea

[0471] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2,5-dimethylaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(4-methyl-1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (27 mg, yield 59%).

[0472] $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 1.65 (d, $J = 6.8$ Hz, 3H), 2.11 (s, 3H), 2.27 (s, 3H), 2.40 (d, $J = 1.0$ Hz, 1H), 4.08 (s, 3H), 4.09 (s, 3H), 5.25 - 5.34 (m, 1H), 6.08 (br, 1H), 6.42 (d, $J = 5.9$ Hz, 1H), 6.80 (s, 1H), 6.93 (s, 1H), 6.96 (br, 1H), 7.61 (s, 1H), 7.68 - 7.73 (m, 2H), 8.41 (d, $J = 5.9$ Hz, 1H)

[0473] Mass spectrometric value (ESI-MS, m/z): 493 ($M^+ + 1$)

Example 94: N-{4-[(6,7-Dimethoxy-4-quinolyl)oxy]phenyl}-N'-[1-(4,5-dimethyl-1,3-thiazol-2-yl)ethyl]urea

[0474] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]aniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(4,5-dimethyl-1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (19 mg, yield 40%).

[0475] $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 1.60 (d, $J = 6.8$ Hz, 3H), 2.28 (s, 3H), 2.31 (s, 3H), 4.03 (s, 3H), 4.04 (s, 3H), 5.19 - 5.26 (m, 1H), 6.29 (br, 1H), 6.44 (d, $J = 5.4$ Hz, 1H), 7.05 (d, $J = 8.8$ Hz, 2H), 7.46 (d, $J = 9.0$ Hz, 2H), 7.48 (s, 1H), 7.55 (s, 1H), 8.04 (br, 1H), 8.43 (d, $J = 5.4$ Hz, 1H)

[0476] Mass spectrometric value (ESI-MS, m/z): 479 ($M^+ + 1$)

Example 95: N-{4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-fluorophenyl}-N'-[1-(4,5-dimethyl-1,3-thiazol-2-yl)ethyl]urea

[0477] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-fluoroaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(4,5-dimethyl-1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (18 mg, yield 38%).

[0478] $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 1.62 (d, $J = 6.8$ Hz, 3H), 2.30 (s, 3H), 2.32 (s, 3H), 4.03 (s, 3H), 4.05 (s, 3H), 5.19 - 5.25 (m, 1H), 6.50 (d, $J = 5.4$ Hz, 1H), 6.58 (br, 1H), 6.87 - 7.05 (m, 2H), 7.51 (s, 1H), 7.71 (br, 1H), 8.22 (d, $J = 9.0$ Hz, 1H), 8.48 (d, $J = 5.4$ Hz, 1H)

[0479] Mass spectrometric value (ESI-MS, m/z): 497 ($M^+ + 1$)

Example 96: N-{4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methylphenyl}-N'-[1-(4,5-dimethyl-1,3-thiazol-2-yl)ethyl]urea

[0480] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methylaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(4,5-dimethyl-1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (18 mg, yield 39%).

[0481] $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 1.60 (d, $J = 6.8$ Hz, 3H), 2.23 (s, 3H), 2.25 (s, 3H), 2.31 (s, 3H), 4.03 (s, 3H), 4.05 (s, 3H), 5.19 - 5.27 (m, 1H), 6.24 (br, 1H), 6.50 (d, $J = 5.4$ Hz, 1H), 6.99 (d, $J = 2.7$ Hz, 1H), 7.01 (s, 1H), 7.10 (s,

1H), 7.47 (s, 1H), 7.54 (s, 1H), 7.75 (d, J = 9.5 Hz, 1H), 8.47 (d, J = 5.6 Hz, 1H)

[0482] Mass spectrometric value (ESI-MS, m/z): 493 (M⁺+1)

Example 97: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-(trifluoromethyl)phenyl]-N'-[1-(4,5-dimethyl-1,3-thiazol-2-yl)ethyl]urea

[0483] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-(trifluoromethyl)aniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(4,5-dimethyl-1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (14 mg, yield 31%).

[0484] ¹H-NMR (CDCl₃, 400 MHz): δ 1.62 (d, J = 6.8 Hz, 3H), 2.29 (s, 3H), 2.32 (s, 3H), 4.04 (s, 3H), 4.06 (s, 3H), 5.21 - 5.25 (m, 1H), 6.49 (d, J = 5.4 Hz, 1H), 6.65 (s, 1H), 7.15 (s, 1H), 7.33 (d, J = 8.8 Hz, 1H), 7.40 (d, J = 2.2 Hz, 1H), 7.52 (s, 1H), 7.52 (s, 1H), 8.20 (dd, J = 9.0 Hz, 1H), 8.51 (d, J = 5.4 Hz, 1H)

[0485] Mass spectrometric value (ESI-MS, m/z): 547 (M⁺+1)

Example 98: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2,5-dimethylphenyl]-N'-[1-(4,5-dimethyl-1,3-thiazol-2-yl)ethyl]urea

[0486] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2,5-dimethylaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(4,5-dimethyl-1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (14 mg, yield 29%).

[0487] ¹H-NMR (CDCl₃, 400 MHz): δ 1.61 (d, J = 6.8 Hz, 3H), 2.12 (s, 3H), 2.21 (s, 3H), 2.26 (s, 3H), 2.29 (s, 3H), 4.05 (s, 3H), 4.06 (s, 3H), 5.20 - 5.26 (m, 1H), 6.13 (s, 1H), 6.36 (d, J = 5.4 Hz, 1H), 6.91 (s, 1H), 6.91 (s, 1H), 7.55 (s, 1H), 7.59 (s, 1H), 7.66 (s, 1H), 8.43 (d, J = 5.4 Hz, 1H)

[0488] Mass spectrometric value (ESI-MS, m/z): 507 (M⁺+1)

Example 99: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methoxyphenyl]-N'-[1-(4,5-dimethyl-1,3-thiazol-2-yl)ethyl]urea

[0489] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methoxyaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(4,5-dimethyl-1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (45 mg, yield 100%).

[0490] ¹H-NMR (CDCl₃, 400 MHz): δ 1.61 (d, J = 6.8 Hz, 3H), 2.31 (s, 3H), 2.33 (s, 3H), 3.66 (s, 3H), 4.01 (s, 3H), 4.04 (s, 3H), 5.26 - 5.29 (m, 1H), 6.46 (d, J = 5.4 Hz, 1H), 6.62 (d, J = 2.4 Hz, 1H), 6.73 (dd, J = 2.4, 8.8 Hz, 1H), 6.76 (s, 1H), 6.78 (s, 1H), 6.76 (s, 1H), 7.41 (s, 1H), 7.46 (s, 1H), 7.54 (s, 1H), 8.20 (d, J = 8.8 Hz, 1H), 8.46 (d, J = 5.4 Hz, 1H)

[0491] Mass spectrometric value (ESI-MS, m/z): 509 (M⁺+1)

Example 100: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]phenyl]-N'-[1-(5-methyl-1,3-thiazol-2-yl)ethyl]urea

[0492] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]aniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(5-methyl-1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (14 mg, yield 31%).

[0493] ¹H-NMR (CDCl₃, 400 MHz): δ 1.49 (d, J = 6.8 Hz, 3H), 2.39 (d, J = 1.2 Hz, 3H), 3.92 (s, 3H), 3.93 (s, 3H), 5.00 - 5.06 (m, 1H), 6.40 (d, J = 5.1 Hz, 1H), 6.87 (d, J = 7.8 Hz, 1H), 7.15 (d, J = 8.8 Hz, 2H), 7.37 (s, 1H), 7.38 (d, J = 1.2 Hz, 1H), 7.50 (s, 1H), 7.52 (d, J = 8.8 Hz, 2H), 8.44 (d, J = 5.1 Hz, 1H), 8.70 (s, 1H)

[0494] Mass spectrometric value (ESI-MS, m/z): 465 (M⁺+1)

Example 101: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-fluorophenyl]-N'-[1-(5-methyl-1,3-thiazol-2-yl)ethyl]urea

[0495] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-fluoroaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(5-methyl-1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (13 mg, yield 29%).

[0496] ¹H-NMR (CDCl₃, 400 MHz): δ 1.66 (d, J = 7.1 Hz, 3H), 2.42 (d, J = 0.7 Hz, 3H), 4.07 (s, 3H), 4.10 (s, 3H), 5.24 - 5.28 (m, 1H), 6.57 (br, 1H), 6.60 (d, J = 5.9 Hz, 1H), 6.90 (dd, J = 2.7, 10.7 Hz, 1H), 6.96 - 6.98 (m, 1H), 7.32 (d, J = 1.2 Hz, 1H), 7.57 (s, 1H), 7.75 (br, 1H), 7.78 (s, 1H), 8.33 (t, J = 8.8 Hz, 1H), 8.48 (d, J = 6.1 Hz, 1H)

[0497] Mass spectrometric value (ESI-MS, m/z): 483 (M⁺+1)

Example 102: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-3-fluorophenyl]-N'-[1-(5-methyl-1,3-thiazol-2-yl)ethyl]urea

[0498] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-3-fluoroaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(5-methyl-1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (16 mg, yield 35%).

[0499] ¹H-NMR (CDCl₃, 400 MHz): δ 1.63 (d, J = 7.1 Hz, 3H), 2.41 (s, 3H), 4.04 (s, 3H), 4.06 (s, 3H), 5.23 - 5.27 (m, 1H), 6.47 (d, J = 5.6 Hz, 1H), 6.54 (br, 1H), 7.08 (t, J = 8.5 Hz, 1H), 7.17 (d, J = 8.5 Hz, 1H), 7.29 (s, 1H), 7.59 - 7.63 (m, 3H), 8.42 (d, J = 5.9 Hz, 1H), 8.57 (s, 1H)

[0500] Mass spectrometric value (ESI-MS, m/z): 483 (M⁺+1)

Example 103: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methylphenyl]-N'-[1-(5-methyl-1,3-thiazol-2-yl)ethyl]urea

[0501] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methylaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(5-methyl-1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (14 mg, yield 31%).

[0502] ¹H-NMR (CDCl₃, 400 MHz): δ 1.63 (d, J = 7.1 Hz, 3H), 2.30 (s, 3H), 2.42 (s, 3H), 4.06 (s, 3H), 4.07 (s, 3H), 5.25 - 5.28 (m, 1H), 6.20 (s, 1H), 6.56 (d, J = 5.9 Hz, 1H), 7.00 (s, 1H), 7.01 (d, J = 2.9 Hz, 1H), 7.17 (s, 1H), 7.28 (s, 1H), 7.57 (s, 1H), 7.64 (s, 1H), 7.79 (d, J = 9.5 Hz, 1H), 8.44 (d, J = 5.9 Hz, 1H)

[0503] Mass spectrometric value (ESI-MS, m/z): 479 (M⁺+1)

Example 104: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-(trifluoromethyl)phenyl]-N'-[1-(5-methyl-1,3-thiazol-2-yl)ethyl]urea

[0504] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-(trifluoromethyl)aniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(5-methyl-1,3-thiazol-2-yl)-1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (12 mg, yield 26%).

[0505] ¹H-NMR (CDCl₃, 400 MHz): δ 1.65 (d, J = 6.8 Hz, 3H), 2.44 (d, J = 0.7 Hz, 3H), 4.07 (s, 3H), 4.09 (s, 3H), 5.23 - 5.30 (m, 1H), 6.37 (d, J = 6.3 Hz, 1H), 6.56 (d, J = 5.6 Hz, 1H), 7.18 (s, 1H), 7.33 (d, J = 1.2 Hz, 1H), 7.36 (dd, J = 2.7, 9.0 Hz, 1H), 7.43 (d, J = 2.7 Hz, 1H), 7.55 (s, 1H), 7.71 (s, 1H), 8.22 (d, J = 9.0 Hz, 1H), 8.51 (d, J = 5.9 Hz, 1H)

[0506] Mass spectrometric value (ESI-MS, m/z): 533 (M⁺+1)

Example 105: N-[4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methoxyphenyl]-N'-[1-(5-methyl-1,3-thiazol-2-yl)ethyl]urea

[0507] 4-[(6,7-Dimethoxy-4-quinolyl)oxy]-2-methoxyaniline (30 mg) was dissolved in chloroform (1 ml) and triethylamine (0.1 ml) to prepare a solution. A solution of triphosgene (35 mg) in chloroform (0.2 ml) was then added to the solution, and the mixture was stirred at room temperature for one hr. Next, a solution of 1-(5-methyl-1,3-thiazol-2-yl)-






1-ethylamine (35 mg) in chloroform (0.2 ml) was added thereto, and the mixture was stirred at room temperature for 10 hr. The stirred mixture was purified by chromatography on silica gel using chloroform/methanol for development to give the title compound (38 mg, yield 83%).






[0508] $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 1.63 (d, $J = 6.8$ Hz, 3H), 2.44 (s, 3H), 3.70 (s, 3H), 4.03 (s, 3H), 4.04 (s, 3H), 5.28 - 5.35 (m, 1H), 6.47 (d, $J = 5.4$ Hz, 1H), 6.49 (s, 1H), 6.76 (dd, $J = 2.2, 8.8$ Hz, 1H), 7.35 (s, 1H), 7.37 (s, 1H), 7.46 (s, 1H), 7.56 (s, 1H), 8.21 (d, $J = 8.8$ Hz, 1H), 8.47 (d, $J = 5.4$ Hz, 1H)






[0509] Mass spectrometric value (ESI-MS, m/z): 495 ($M^+ + 1$)







[0510] Compounds of Examples 1 to 38 and 41 to 105 had the following respective chemical structures and c-fms autophosphorylation 50% inhibitory concentrations (IC_{50}).






Table 1







Example	Chemical structure	IC ₅₀ (μm)
1		0.0024
2		0.0022
3		0.0024
4		0.0040
5		0.0022







5 10	6		0.0060
15 20	7		0.0202
25 30	8		0.0225
35 40	9		0.0174
45 50	10		0.0071







5 11		0.0117
10 12		0.0109
15 20 25 13		0.0061
30 35 40 14		0.0130
45 50 15		0.0057






5 10	16		0.0020
15 20	17		0.0039
25 30	18		0.0037
35 40	19		0.0081
45 50	20		0.0062
55	21		0.0098






22		0.0077
23		0.0102
24		0.0132
25		0.0219
26		0.0163






27		0.0076
28		0.0065
29		0.0023
30		0.0031
31		0.0129
32		0.0035






33		0.0065
34		0.0063
35		0.0121
36		0.0311
37		0.0050
38		0.0057






5 41		0.0028
10 42		0.0385
15 43		0.0031
20 44		0.0018
25 45		0.0222
30 46		0.0073






47		0.0038
48		0.0400
49		0.0032
50		0.0072
51		0.0076






5 10 15 20 25 30 35 40 45 50 55	52		0.0055
	53		0.0027
	54		0.0695
	55		0.0454
	56		0.1933





57		0.1275
58		0.0809
59		0.0203
60		0.1152
61		0.0443






5 10	62		0.0294
15 20	63		0.0309
25 30	64		0.0094
35 40	65		0.0139
45 50 55	66		0.0147





5 67		0.0057
10 68		0.0098
15 69		0.0115
20 70		0.0036
25 71		0.0030






72		0.0049
73		0.0030
74		0.0050
75		0.0021
76		0.0071






5 10	77		0.0047
15 20	78		0.0029
25 30	79		0.0069
35 40	80		0.0062
45 50 55	81		0.0062


5 10	82		0.0089
15 20	83		0.0102
25 30	84		0.0102
35 40 45	85		0.0174

5 10	86		0.0125
15 20	87		0.0265
25 30	88		0.0029
35 40	89		0.0031
45 50 55	90		0.0036

5 10	91		0.0032
15 20	92		0.0084
25 30	93		0.0386
35 40 45	94		0.0535

5 10	95		0.0383
15 20	96		0.1468
25 30	97		0.1202
35 40	98		0.4260
45 50	99		0.0126

100		0.0157
101		0.0201
102		0.0203
103		0.0696
104		0.0220

105		0.0127
-----	--	--------

Pharmacological Test Example 1: Measurement of inhibitory activity against c-fms tyrosine kinase activity within macrophage-like cell line BAC-1.2F5 by M-CSF stimulation

[0511] Macrophage-like cell line BAC-1.2F5 which expresses c-fms as a M-CSF receptor on cell membranes (CLAUDIA M. et al., J. Cell. Physiol., 130, 420-427 (1987), "Isolation and Characterization of a Cloned Growth Factor Dependent Macrophage Cell Line, BAC1.2F5") was cultured at 37°C in a DMEM medium containing L-cell supernatant and 10% fetal calf serum (GIBCO BRL) in a 5% CO₂ incubator until about 50 to 70% confluent. The harvested cells were inoculated into wells, each containing the same medium, in a collagen type I coat 96-well flat-bottom plate in an amount of 5.0 x 10⁴ per well, and cultured at 37°C for 12 hr. Thereafter, the medium was replaced with a DMEM medium containing a 0.1% fetal calf serum and cultured for 12 hr. Thereafter, a test compound dissolved in dimethyl sulfoxide was added to each well, and cells were incubated at 37°C for one hr. Human recombinant M-CSF was added to a final concentration of 50 ng/ml, and cells were stimulated at 37°C for 6 min. After the medium was removed, 50 µl of a solubilization buffer (20 mM HEPES (pH 7.4), 150 mM NaCl, 0.2% Triton X-100, 10% glycerol, 5 mM sodium orthovanadate, 5 mM disodium ethylenediaminetetraacetate, and 2 mM Na₄P₂O₇) was added thereto. The mixture was shaken at 4°C for 2 hr to prepare a cell extract as an assay sample which was then used for an ELISA assay.

[0512] The plate for the ELISA assay was prepared as follows. Phosphate buffered saline (50 µl, pH 7.4) containing 5 µg/ml of anti-phospho-tyrosine antibody (PY20; Transduction Laboratories) was added to a microplate for ELISA (Maxisorp; purchased from NUNC), followed by standing at 4°C overnight to be immobilized on the wells. Thereafter, each well was washed twice with 200 µl of a Ca- and Mg-free phosphate buffer containing 0.1% Tween-20 (hereinafter referred to as "PBS-T"), and 200 µl of a blocking solution (Dainippon Pharmaceutical Co. Ltd.: BlockAce) was added to each well, and the wells were then allowed to stand at room temperature for 2 hr for blocking.

[0513] Separately, phosphate buffered saline (50 µl, pH 7.4) containing 5 µg/ml of anti-phospho-tyrosine antibody (PY20; Transduction Laboratories) was added to a microplate for ELISA (Maxisorp; purchased from NUNC), followed by standing at 4°C overnight to form a solid phase on the wells. After washing of the plate, 300 µl of a blocking solution was added, followed by standing at room temperature for 2 hr to perform blocking. After washing, the whole quantity of the cell extract was transferred to the wells, and the plate was then allowed to stand at 4°C overnight. After washing, anti-c-fms antiserum (Upstate Biotechnology, Inc.) was allowed to react at room temperature for one hr, and, after washing, a peroxidase-labeled anti-rabbit Ig antibody (Amersham) was allowed to react at room temperature for one hr. After washing, a chromophoric substrate for peroxidase (Sumitomo Bakelite Co., Ltd.) was added thereto to initiate a reaction. After a suitable level of color development, a reaction termination solution was added to stop the reaction, and the absorbance at 450 nm was measured with a microplate reader. The c-fms phosphorylation activity for each well was determined by presuming the absorbance without the addition of the test compound and with the addition of M-CSF to be 100% c-fms phosphorylation activity and the absorbance without the addition of the test compound and M-CSF to be 0% c-fms phosphorylation activity. The concentration of the test compound was varied on several levels, the c-fms phosphorylation inhibitory rate was determined for each case, and the concentration of the test compound necessary for inhibiting 50% of c-fms phosphorylation (IC₅₀) was calculated. The results were as shown in the table described before.

Pharmacological Test Example 2: Measurement of inhibitory activity against M-CSF-dependent cell growth of macrophage-like cell line BAC-1.2F5

[0514] Macrophage-like cell line BAC-1.2F5 which expresses c-fms as a M-CSF receptor on cell membranes (CLAUDIA M. et al., J. Cell. Physiol., 130, 420-427 (1987), "Isolation and Characterization of a Cloned Growth Factor De-

pendent Macrophage Cell Line, BAC1.2F5") was cultured at 37°C in a DMEM medium containing L-cell supernatant and 10% fetal calf serum (purchased from GIBCO BRL) in a 5% CO₂ incubator until about 50 to 70% confluent. The harvested cells were inoculated into wells, each containing a 10% fetal calf serum-containing DMEM medium with human recombinant M-CSF being added thereto to a final concentration of 50 ng/ml, in a 96-well flat-bottom plate at a density of 5.0×10^3 per well, and cultured at 37°C for 24 hr. A test compound dissolved in dimethyl sulfoxide was then added to each well to a target concentration, and cells were incubated at 37°C in a 5% CO₂ incubator for 72 hr. Thereafter, 20 µl of an MTS reagent (purchased from Promega) was added to each well, followed by incubation at 37°C in a 5% CO₂ incubator for 2 hr. Thereafter, absorbance at 490 nm was measured with a microplate reader. The cell growth rate with the addition of the test compound was determined by presuming the absorbance without the addition of the test compound and with the addition of M-CSF to be 100% cell growth and the absorbance without the addition of the test compound and M-CSF to be 0% cell growth. The concentration of the test compound was varied on several levels, and the cell growth rate in each case was calculated. As a result, a group of compounds according to the present invention inhibited M-CSF-dependent cell growth of BAC-1.2F5 cell line in a concentration-dependent manner. A representative example thereof is shown in Fig. 1.

Pharmacological Test Example 3: Osteoclast differentiation-inducing test by cocultivation of mouse bone marrow cells and cell line for osteoblasts

[0515] Mouse osteoblast-like cell line KS483 (YAMASHITA T. et al., Bone, 19, 429-436 (1996), "Subcloning of Three Osteoblastic Cell with Distinct Differentiation Phenotypes from the Mouse Osteoblastic Cell line KS-4") was cultured in an α -MEM medium containing a 10% fetal calf serum (purchased from GIBCO BRL) until about 90% confluent. The cells were seeded into wells, each containing the same medium containing a 10% fetal calf serum, in a 48 well flat-bottom plate at a density of 2.0×10^3 per well. Bone marrow cells harvested from femorotibialis of 4-week old male DDY mice (purchased from Japan SLC, Inc.) were seeded into the same wells in which KS483 was cultured at a density of 2.0×10^4 per well, and 1,25(OH)₂VD₃ was added to a final concentration of 10^{-8} mol/L. Further, the test compound dissolved in dimethyl sulfoxide was added to each well to a target concentration, followed by incubation in a 5% CO₂ incubator at 37°C for 6 days. The medium was replaced every two days, and each time the replacement was carried out, 1,25(OH)₂VD₃ and the test compound were added to the above concentration. Staining (ACID PHOSPHATASE LEUKOCYTE; SIGMA) and observation were carried out after cultivation for 6 days using tartaric acid-resistant acid phosphatase (hereinafter referred to as "TRAP") activity (a specific marker found on the surface of osteoclast cell) as an index. After staining, observation under an optical microscope was carried out. As a result, it was found that a group of compounds according to the present invention reduced the number of osteoclasts stained by TRAP activity in a concentration-dependent manner. At the same time, the TRAP activity was evaluated by the measurement of enzyme activity using p-nitrophenylphosphate (SIGMA104 phosphatase substrate; SIGMA; hereinafter referred to as "PNP"). The measurement of enzyme activity was carried out with PNP as follows. The medium in each well was removed after 6 days culture, and cells were fixed with an acetone-water-citrate buffer (weight ratio = 15 : 9 : 1) mixed solution. After washing, PNP was dissolved in a water-acetate buffer-tartrate buffer (weight ratio = 22 : 1 : 1) mixed solution to a final concentration of 0.4 mg/ml. This PNP solution was added to each well and was stirred at room temperature for one hr. After stirring, 50 µl of 0.5 N NaOH was further added, and 100 µl of the reaction solution in each well was dispensed in a 96 well flat-bottom plate, and absorbance at 405 nm was then measured with a microplate reader. For each well, TRAP activity was measured by presuming the absorbance in the case of culture without the addition of the test compound and with the addition of only KS483 and mouse bone marrow cells in the presence of 1,25(OH)₂VD₃ to 100% TRAP activity and the absorbance without the addition of the test compound and the mouse bone marrow cells to 0% TRAP activity. The concentration of the test compound was varied on several levels, and, for each case, the rate of inhibition against TRAP activity was determined. As a result, a group of compounds according to the present invention reduced TRAP activity in a concentration-dependent manner. That is, the differentiation of osteoclasts was inhibited in a dose-dependent manner. A typical example thereof is shown in Fig. 2.

Pharmacological Test Example 4: Inhibitory effect in nude mouse bone metastasis model using human melanoma cells (A375)

[0516] An A375 human melanoma cell line (obtained from Japanese Foundation for Cancer Research) was inoculated under anesthesia with Ketamine-Medetomidine into the left ventricle of nude mice to 1.0×10^5 cells/head, and, on the day of transplantation, the mice were grouped so that the groups each consisted of five mice and had an identical body weight. From the next day of the transplantation, for 28 days twice a day, the test compound was orally administered to the treatment group at 40 mg/kg/day, and the medium was orally administered to the control group. On 21 and 28 days after the transplantation, the mice were radiographed. For the images thus obtained, the bone resorption pore area in femora and fibiae attributable to bone metastasis of melanoma in each individual were calculated with an

image analysis software (Image Gauge, FUJI FILM) to determine the average value for each group. The drug efficacy was evaluated by comparing the area of the test compound administered group with the area of the control group. As a result, a group of compounds according to the present invention were found to inhibit the expansion of the area of bone resorption attributable to bone metastatic melanoma cells. A representative example thereof is shown in Fig. 3.

Pharmacological Test Example 5: Inhibitory effect in nude rat bone metastasis model using human melanoma cells (A375)

[0517] An A375 human melanoma cell line (obtained from Japanese Foundation for Cancer Research) was inoculated under anesthesia with Ketamine-Medetomidine into the left ventricle of nude mice to 5.0×10^5 cells/head, and, on the day of transplantation, the mice were grouped so that the groups each consisted of eight mice and had an identical body weight. From the next day of the transplantation, for 21 days once a day, the test compound was orally administered to the treatment group at 50 mg/kg, and the vehicle was orally administered to the control group. On 21 day after the transplantation, the mice were radiographed. For the images thus obtained, the femorotibial bone resorption pore area attributable to bone metastasis of melanoma in each individual were calculated with an image analysis software (Image Gauge, FUJI FILM) to determine the average value for each group. The drug efficacy was evaluated by comparing the area of the test compound administered group with the area of the control group. As a result, a group of compounds according to the present invention were found to inhibit the expansion of the area of bone resorption attributable to bone metastatic melanoma cells. A representative example thereof is shown in Fig. 4.

Pharmacological Test Example 6: Effect of inhibiting increase in number of osteoclasts in spayed model rats

[0518] Six-week old female SD rats (purchased from Japan SLC, Inc.) were grouped so that the groups each consisted of six rats and had an identical body weight. On the 7th day after grouping, both ovaries were spayed to prepare an osteoporosis model. For 28 days from the day of grouping once a day, the test compound was orally administered at 20 mg/kg/day, and the vehicle was orally administered to the control group. On the 21st day after the transplantation, tibias of both feet were harvested and were fixed with 10% buffered formalin. Sections were then prepared, and TRAP staining was carried out. An optical microscope was interlocked with a personal computer, and the number of TRAP-positive osteoclasts in a primary cancellous zone of tibial metaphyseal part was measured with a bone form analysis software (Luzex F Bone System, NIRECO CORPORATION). TRAP-positive cells having two or more nuclei were regarded as osteoclasts. In the measurement, a line connecting one end of the growth cartilage to a metaphyseal junction (hereinafter referred to as "GCMJ") or a line which is parallel to a line connecting both ends of the growth cartilage in its lower part and is in contact with GCMJ is provided as a base line. A rectangular region showing an area of 0.96 mm^2 defined by about 0.4 mm from the base line toward shaft and, in the lateral direction, about 2.4 mm distant from GCMJ as the center was set as a measurement viewing field. As a result, a group of compounds according to the present invention significantly reduced the number of osteoclasts. A typical example thereof is shown in Fig. 5.

Pharmacological Test Example 7: Measurement of inhibitory activity against KDR phosphorylation using ELISA method

[0519] Human KDR-transfected NIH3T3 cells (Sawano A et al., Cell Growth & Differentiation, 7, 213-221 (1996)) were cultured in a DMEM medium containing 10% fetal calf serum (purchased from GIBCO BRL) within a 5% carbon dioxide incubator until 50 to 70% confluent. The harvested cells were seeded into wells, containing the same medium, in a collagen type 1 coat 96-well flat-bottom plate in an amount of 1.5×10^4 per well, and cultured at 37°C overnight. The medium was replaced with a 0.1% fetal calf serum-containing DMEM medium. The test compound dissolved in dimethyl sulfoxide was added to each well, and the cultivation was continued at 37°C for additional one hr. A human recombinant vascular endothelial growth factor (hereinafter abbreviated to "VEGF") was added to a final concentration of 100 ng/ml, followed by stimulation of cells at 37°C for 2 min. The medium was removed, and the cells were washed with phosphate buffered saline (pH 7.4). A solubilization buffer (50 μl) (20 mM HEPES (pH 7.4), 150 mM NaCl, 0.2% Triton X-100, 10% glycerol, 5 mM sodium orthovanadate, 5 mM disodium ethylenediaminetetraacetate, and 2 mM $\text{Na}_4\text{P}_2\text{O}_7$) was then added thereto. The mixture was shaken at 4°C for 2 hr to prepare a cell extract.

[0520] Separately, phosphate buffered saline (50 μl , pH 7.4) containing 5 $\mu\text{g/ml}$ of anti-phospho-tyrosine antibody (PY20; purchased from Transduction Laboratories) was added to a microplate for ELISA (Maxisorp; purchased from NUNC), followed by standing at 4°C overnight to be immobilized on the wells. After washing of the plate, 300 μl of a blocking solution was added, followed by standing at room temperature for 2 hr to perform blocking. After washing, the whole cell extract was transferred and was allowed to stand at 4°C overnight. After washing, an anti-KDR antibody (purchased from Santa Cruz Biotechnology) was allowed to react at room temperature for one hr. Further, after washing, a peroxidase-labeled anti-rabbit Ig antibody (purchased from Amersham) was allowed to react at room temperature for one hr. After washing, a chromophoric substrate for peroxidase (purchased from Sumitomo Bakelite Co., Ltd.) was

added thereto to initiate a reaction. After a suitable level of color development, a reaction termination solution was added to stop the reaction, and the absorbance at 450 nm was measured with a microplate reader. The KDR phosphorylation activity for each well was determined by presuming the absorbance without the addition of the test compound and with the addition of VEGF to be 100% KDR phosphorylation activity and the absorbance without the addition of the test compound and VEGF to be 0% KDR phosphorylation activity. The concentration of the test compound was varied on several levels, the KDR phosphorylation inhibitory rate was determined for each case, and the concentration of the test compound necessary for inhibiting 50% of KDR phosphorylation (IC_{50}) was calculated.

[0521] KDR phosphorylation 50% inhibitory concentration (IC_{50}) regarding typical examples of a group of compounds according to the present invention was as shown in Table 2. The selection ratio of the KDR phosphorylation inhibitory activity to the c-fms phosphorylation inhibitory activity is also shown in Table 2. When KDR phosphorylation 50% inhibitory concentration [μM]/c-fms phosphorylation 50% inhibitory concentration [μM] was less than 5, the selection ratio was indicated as "0"; when the value was 5 or more, the selection ratio was indicated as "1"; when the value was 10 or more, the selection ratio was indicated as "2"; and when the value was 50 or more, the selection ratio was indicated as "3."

Table 2

Ex. No.	KDR (μM)	Selection ratio
1	0.0011	0
2	0.0323	2
3	< 0.0010	0
4	0.0010	0
5	0.0214	1
6	0.0018	0
7	0.0031	0
8	0.3897	2
9	0.0027	0
10	0.0030	0
11	0.1232	2
12	0.0028	0
13	< 0.0010	0
14	0.0688	1
15	< 0.0010	0
16	0.0031	0
17	0.3149	3
18	0.0018	0
19	0.0015	0
20	0.1462	2
21	0.0021	0
22	0.0039	0
23	0.6000	3
24	0.0024	0
25	0.0035	0
26	0.1550	1
27	0.0020	0
28	< 0.0010	0

EP 1 535 910 A1

Table 2 (continued)

Ex. No.	KDR (μM)	Selection ratio
29	0.0273	2
30	< 0.0010	0
31	0.0010	0
32	0.0294	1
33	< 0.0010	0
34	0.2963	2
35	0.3097	2
36	0.1343	0
41	0.0188	1
42	0.0166	0
45	0.0025	0
46	0.5071	3
47	0.3097	3
48	0.0027	0
49	0.0454	2
50	0.0283	0
51	0.0382	1
52	0.0072	0
53	0.0374	2
54	0.5740	1
55	0.0016	0
56	0.0096	0
57	0.0036	0
58	0.0021	0
59	0.0087	0
60	0.0019	0
61	0.0192	0
62	0.0027	0
63	0.0026	0
64	0.0098	0
65	0.0319	0
66	0.0165	0
67	0.0029	0
68	0.0056	0
69	0.0052	0
70	0.0551	2
71	0.0399	2
72	0.0356	1


Table 2 (continued)

Ex. No.	KDR (μ M)	Selection ratio
73	0.0433	2
74	0.3176	3
75	0.1463	3
76	0.4368	3
77	0.1012	2
78	0.0539	2
79	0.0952	2
80	0.5257	3
81	1.0521	3
82	0.6501	3
83	0.0079	0
84	0.6609	3
85	0.0216	0
86	0.2081	2
87	0.2731	2
88	0.0087	0
89	0.0174	1
90	0.1110	2
91	0.0288	1
92	0.0043	0
93	0.2680	1
99	0.2924	2
100	0.3108	2
101	0.2387	2
102	0.2193	2
103	0.0909	0
104	0.0361	0
105	0.3096	2

[0522] Thus, the compounds according to the present invention have high c-fms phosphorylation inhibitory activity in vitro and selectively inhibit c-fms phosphorylation rather than KDR phosphorylation in vitro.

Claims

1. A compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof:



wherein

X represents CH or N;

Z represents O or S;

R¹, R², and R³, which may be the same or different, represent a hydrogen atom; a halogen atom; hydroxyl; cyano; C₁₋₆ alkyl; C₁₋₆ alkoxy; C₂₋₆ alkenyl; C₂₋₆ alkynyl; nitro; -NR¹⁰⁶R¹⁰⁷ wherein R¹⁰⁶ and R¹⁰⁷, which may be the same or different, represent a hydrogen atom or C₁₋₄ alkyl in which the alkyl group is optionally substituted by hydroxyl, -OR¹⁰⁸ wherein R¹⁰⁸ represents C₁₋₄ alkyl, or -NR¹⁰⁹R¹¹⁰ wherein R¹⁰⁹ and R¹¹⁰, which may be the same or different, represent a hydrogen atom or C₁₋₄ alkyl; -CONR¹¹¹R¹¹² wherein R¹¹¹ and R¹¹², which may be the same or different, represent a hydrogen atom or C₁₋₄ alkyl in which the alkyl group is optionally substituted by hydroxyl, -OR¹¹³ wherein R¹¹³ represents C₁₋₄ alkyl, or -NR¹¹⁴R¹¹⁵ wherein R¹¹⁴ and R¹¹⁵, which may be the same or different, represent a hydrogen atom or C₁₋₄ alkyl; or -COOR¹¹⁶ wherein R¹¹⁶ represents a hydrogen atom or C₁₋₄ alkyl in which the alkyl group is optionally substituted by hydroxyl, -OR¹¹⁷ wherein R¹¹⁷ represents C₁₋₄ alkyl, or -NR¹¹⁸R¹¹⁹ wherein R¹¹⁸ and R¹¹⁹, which may be the same or different, represent a hydrogen atom or C₁₋₄ alkyl in which the C₁₋₆ alkyl, C₁₋₆ alkoxy, C₂₋₆ alkenyl, and C₂₋₆ alkynyl groups are optionally substituted by a halogen atom; hydroxyl; C₁₋₄ alkyl; C₁₋₄ alkoxy; C₁₋₄ alkoxycarbonyl; amino in which one or two hydrogen atoms on the amino group each are optionally substituted by C₁₋₄ alkyl optionally substituted by hydroxyl or C₁₋₄ alkoxy; group R¹⁵R¹⁶N-C(=O)-O- wherein R¹⁵ and R¹⁶, which may be the same or different, represent a hydrogen atom or C₁₋₄ alkyl in which the alkyl group is optionally substituted by hydroxyl or C₁₋₄ alkoxy; or group R¹⁷-(S)_m- wherein R¹⁷ represents a saturated or unsaturated three- to seven-membered carbocyclic or heterocyclic group optionally substituted by a halogen atom or C₁₋₄ alkyl and m is 0 (zero) or 1,

R⁴ represents a hydrogen atom,

R⁵, R⁶, R⁷, and R⁸, which may be the same or different, represent a hydrogen atom, a halogen atom, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio, trifluoromethyl, nitro, or amino,

R⁹ and R¹⁰, which may be the same or different, represent a hydrogen atom, C₁₋₆ alkyl, or C₁₋₄ alkylcarbonyl, and any one of R¹¹ and R¹² represents a hydrogen atom while the other represents C₁₋₄ alkyl, and R¹³ represents a saturated or unsaturated three- to seven-membered carbocyclic or heterocyclic group or a saturated or unsaturated nine- to twelve-membered bicyclic carbocyclic group in which the carbocyclic and heterocyclic groups are optionally substituted by a halogen atom; hydroxyl; C₁₋₄ alkyl; C₁₋₄ alkoxy; C₁₋₄ alkylthio; trifluoromethyl; nitro; or -NR¹³⁷R¹³⁸ wherein R¹³⁷ and R¹³⁸, which may be the same or different, represent a hydrogen atom or C₁₋₄ alkyl in which the alkyl group is optionally substituted by hydroxyl, -OR¹³⁹ wherein R¹³⁹ represents C₁₋₄ alkyl, or -NR¹⁴⁰R¹⁴¹ wherein R¹⁴⁰ and R¹⁴¹, which may be the same or different, represent a hydrogen atom or C₁₋₄ alkyl, or


R¹¹ represents a hydrogen atom, and R¹² and R¹³ may combine with a carbon atom attached thereto to form a saturated or unsaturated nine- to twelve-membered bicyclic carbocyclic group.

2. The compound according to claim 1, wherein X represents CH.

3. The compound according to claim 1 or 2, wherein Z represents O.

4. The compound according to any one of claims 1 to 3, wherein R¹ and R⁴ represent a hydrogen atom.

5. The compound according to any one of claims 1 to 4, wherein R⁹ and R¹⁰ represent a hydrogen atom.
6. The compound according to any one of claims 1 to 5, wherein R² and R³, which may be the same or different, represent C₁₋₆ alkoxy, said alkoxy group being optionally substituted by a halogen atom; hydroxyl; C₁₋₄ alkyl; C₁₋₄ alkoxy; C₁₋₄ alkoxycarbonyl; amino in which one or two hydrogen atoms on the amino group each are optionally substituted by C₁₋₄ alkyl optionally substituted by hydroxyl or C₁₋₄ alkoxy; or a saturated or unsaturated three- to seven-membered carbocyclic or heterocyclic group.
7. The compound according to any one of claims 1 to 6, wherein at least one of R⁵, R⁶, R⁷ and R⁸ represents a halogen atom, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio, trifluoromethyl, nitro, or amino, and the other(s) represents a hydrogen atom.
8. The compound according to any one of claims 1 to 6, wherein all of R⁵, R⁶, R⁷ and R⁸ represent a hydrogen atom.
9. The compound according to any one of claims 1 to 8, wherein any one of R¹¹ and R¹² represents a hydrogen atom and the other represents C₁₋₄ alkyl, and R¹³ represents phenyl, naphthyl, imidazolyl, oxazolyl, thiazolyl, pyrazolyl, isoxazolyl, or isothiazolyl, said groups being optionally substituted by a halogen atom, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio, trifluoromethyl, nitro, or amino in which one or two hydrogen atoms on the amino group each are optionally substituted by C₁₋₄ alkyl, or
R¹¹ represents a hydrogen atom, and R¹² and R¹³ combine with a carbon atom attached thereto to form 1,2,3,4-tetrahydronaphthalene or indan.
10. The compound according to claim 1, represented by formula (Ia):



wherein

- X represents CH or N,
R¹⁸ and R¹⁹, which may be the same or different, represent C₁₋₆ alkoxy, said alkoxy group being optionally substituted by a halogen atom; hydroxyl; C₁₋₄ alkyl; C₁₋₄ alkoxycarbonyl; amino in which one or two hydrogen atoms on the amino group each are optionally substituted by C₁₋₄ alkyl optionally substituted by hydroxyl or C₁₋₄ alkoxy; or a saturated or unsaturated three- to seven-membered carbocyclic or heterocyclic group,
R²⁰, R²¹, R²², and R²³, which may be the same or different, represent a hydrogen atom, a halogen atom, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio, trifluoromethyl, nitro, or amino,
any one of R²⁴ and R²⁵ represents a hydrogen atom and the other represents C₁₋₄ alkyl, and R²⁶ represents phenyl, naphthyl, imidazolyl, oxazolyl, thiazolyl, pyrazolyl, isoxazolyl, or isothiazolyl, said groups being optionally substituted by a halogen atom, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio, trifluoromethyl, nitro, or amino in which one or two hydrogen atoms on the amino group each are optionally substituted by C₁₋₄ alkyl, or
R²⁴ represents a hydrogen atom, and R²⁵ and R²⁶ combine with a carbon atom attached thereto to form 1,2,3,4-tetrahydronaphthalene or indan.
11. The compound according to claim 10, wherein X represents CH.
12. The compound according to claim 10 or 11, wherein R¹⁸ and R¹⁹, which may be the same or different, represent C₁₋₆ alkoxy optionally substituted by a saturated or unsaturated three- to seven-membered carbocyclic or heterocyclic group.

13. The compound according to any one of claims 10 to 12, wherein at least one of R^{20} , R^{21} , R^{22} and R^{23} represents a halogen atom, C_{1-4} alkyl, C_{1-4} alkoxy, C_{1-4} alkylthio, trifluoromethyl, nitro, or amino, and the other(s) represents a hydrogen atom.

14. The compound according to any one of claims 10 to 12, wherein R^{20} and R^{21} , which may be the same or different, represent a halogen atom, C_{1-4} alkyl, C_{1-4} alkoxy, C_{1-4} alkylthio, trifluoromethyl, nitro, or amino, and R^{22} and R^{23} represent a hydrogen atom.


15. The compound according to any one of claims 10 to 12, wherein R^{21} and R^{22} , which may be the same or different, represent a halogen atom, C_{1-4} alkyl, C_{1-4} alkoxy, C_{1-4} alkylthio, trifluoromethyl, nitro, or amino, and R^{20} and R^{23} represent a hydrogen atom.

16. The compound according to any one of claims 10 to 12, wherein all of R^{20} , R^{21} , R^{22} , and R^{23} represent a hydrogen atom.

17. The compound according to any one of claims 10 to 16, wherein R^{26} represents thiazolyl.

18. The compound according to any one of claims 10 to 16, wherein R^{26} represents 4-fluorophenyl.

19. The compound according to claim 1, represented by formula (Ib)




wherein

R^{31} represents a hydrogen atom, a fluorine atom at 2-position, a fluorine atom at 3-position, methoxy at 2-position, methoxy at 3-position, or methyl at 2- and 5-positions,

R^{32} represents methyl, and


R^{33} represents a hydrogen atom, methyl at 1-position, methyl at 2-position, or methyl at 1- and 2-positions.

20. The compound according to claim 19, wherein the compound represented by formula (Ib) is represented by formula (Ib-1)




wherein R^{31} , R^{32} , and R^{33} are as defined in formula (Ib).

21. The compound according to claim 19, wherein the compound represented by formula (Ib) is represented by formula (Ib-2)



wherein R^{31} , R^{32} , and R^{33} are as defined in formula (Ib).

22. The compound according to claim 1, represented by formula (Ic)




wherein

R^{41} represents a hydrogen atom, a fluorine atom at 2-position, a fluorine atom at 3-position, a chlorine atom at 2-position, a chlorine atom at 3-position, methyl at 2- and 3-positions, methyl at 2- and 5-positions, methoxy at 2-position, methoxy at 3-position, methyl at 2-position, or trifluoromethyl at 2-position,

R^{42} represents methyl,

R^{43} represents a fluorine atom at 4-position, a bromine atom at 3-position, a bromine atom at 4-position, methoxy at 2-position, methoxy at 3-position, methoxy at 4-position, a chlorine atom at 4-position, methyl at 4-position, or nitro at 4-position.

23. The compound according to claim 1, represented by formula (Id)



wherein

X represents CH or N,


R^{51} represents a hydrogen atom, a fluorine atom at 2-position, a fluorine atom at 3-position, methoxy at 2-position, methoxy at 3-position, or methyl at 2- and 5-positions,

R^{52} represents methyl,

R^{53} represents imidazolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, or isothiazolyl in which one or two hydrogen atoms on the groups are optionally substituted by a halogen atom or C_{1-4} alkyl, and

R^{54} and R^{55} , which may be the same or different, represent a hydrogen atom or C_{1-6} alkyl in which the alkyl group is optionally substituted by hydroxyl; a halogen atom; $-OR^{56}$ wherein R^{56} represents C_{1-4} alkyl; $-NR^{57}R^{58}$ wherein R^{57} and R^{58} , which may be the same or different, represent a hydrogen atom or C_{1-4} alkyl in which the alkyl group is optionally substituted by hydroxyl or $-OR^{59}$ wherein R^{59} represents C_{1-4} alkyl; or a saturated or unsaturated three- to seven-membered carbocyclic or heterocyclic group in which the carbocyclic and heterocyclic groups are optionally substituted by one or two halogen atoms or C_{1-4} alkyl.


24. The compound according to claim 23, wherein X represents CH, and R^{52} represents



25. The compound according to claim 24, wherein R⁵⁴ and R⁵⁵ represent methyl.

26. The compound according to claim 24, wherein R⁵⁴ represents methyl, and R⁵⁵ represents C₁₋₄ alkyl substituted by a saturated or unsaturated five- or six-membered carbocyclic or heterocyclic group.


27. The compound according to claim 23, wherein X represents CH, and R⁵² represents



28. The compound according to claim 27, wherein R⁵⁴ and R⁵⁵ represent methyl.

29. The compound according to claim 27, wherein R⁵⁴ represents methyl, and R⁵⁵ represents C₁₋₄ alkyl substituted by a saturated or unsaturated five- or six-membered carbocyclic or heterocyclic group.


30. The compound according to claim 23, wherein X represents N, and R⁵² represents



31. The compound according to claim 30, wherein R⁵⁴ and R⁵⁵ represent methyl.

32. The compound according to claim 30, wherein R⁵⁴ represents methyl, and R⁵⁵ represents C₁₋₄ alkyl substituted by a saturated or unsaturated five- or six-membered carbocyclic or heterocyclic group.


33. The compound according to claim 23, wherein X represents N, and R⁵² represents



34. The compound according to claim 33, wherein R⁵⁴ and R⁵⁵ represent methyl.

35. The compound according to claim 33, wherein R⁵⁴ represents methyl, and R⁵⁵ represents C₁₋₄ alkyl substituted by a saturated or unsaturated five- or six-membered carbocyclic or heterocyclic group.

36. The compound according to claim 1, represented by formula (Ie)



wherein

R^{601} represents a hydrogen atom, a fluorine atom at 2-position, a fluorine atom at 3-position, a chlorine atom at 2-position, a chlorine atom at 3-position, methyl at 2- and 3-positions, methyl at 2- and 5-positions, methoxy at 2-position, methoxy at 3-position, methyl at 2-position, or trifluoromethyl at 2-position,

R^{602} represents methyl,

X represents N or CH,

R^{604} and R^{605} , which may be the same or different, represent a hydrogen atom or C_{1-6} alkyl in which the alkyl group is optionally substituted by hydroxyl; a halogen atom; $-OR^{606}$ wherein R^{606} represents C_{1-4} alkyl; $-NR^{607}R^{608}$ wherein R^{607} and R^{608} , which may be the same or different, represent a hydrogen atom or C_{1-4} alkyl in which the alkyl group is optionally substituted by hydroxyl or $-OR^{609}$ wherein R^{609} represents C_{1-4} alkyl; or a saturated or unsaturated three- to seven-membered carbocyclic or heterocyclic group in which the carbocyclic and heterocyclic groups are optionally substituted by one or two halogen atoms or C_{1-4} alkyl, and

R^{611} , R^{612} , R^{613} , R^{614} , and R^{615} , which may be the same or different, represent a hydrogen atom; C_{1-6} alkyl; $-OR^{616}$ wherein R^{616} represents C_{1-4} alkyl; a halogen atom; nitro; or $-NR^{617}R^{618}$ wherein R^{617} and R^{618} , which may be the same or different, represent a hydrogen atom or C_{1-4} alkyl in which the alkyl group is optionally substituted by hydroxyl, $-OR^{619}$ wherein R^{619} represents C_{1-4} alkyl, or $-NR^{620}R^{621}$ wherein R^{620} and R^{621} , which may be the same or different, represent a hydrogen atom or C_{1-4} alkyl.

37. The compound according to claim 36, wherein X represents CH and all of R^{611} , R^{612} , R^{613} , R^{614} , and R^{615} represent a hydrogen atom, or any one of R^{611} , R^{612} , R^{613} , R^{614} , and R^{615} represents a group other than a hydrogen atom and the remaining groups represent a hydrogen atom.

38. The compound according to claim 37, wherein all of R^{611} , R^{612} , R^{613} , R^{614} , and R^{615} represent a hydrogen atom, or any one of R^{611} , R^{612} , R^{613} , R^{614} , and R^{615} represents C_{1-6} alkyl, $-OR^{616}$, a halogen atom, or nitro and the remaining groups represent a hydrogen atom.

39. The compound according to claim 38, wherein R^{611} represents methoxy and R^{612} , R^{613} , R^{614} , and R^{615} represent a hydrogen atom, or R^{612} represents a bromine atom or methoxy and R^{611} , R^{613} , R^{614} , and R^{615} represent a hydrogen atom, or R^{613} represents a bromine atom, a chlorine atom, a fluorine atom, methyl, methoxy, or nitro and R^{611} , R^{612} , R^{614} , and R^{615} represent a hydrogen atom.

40. The compound according to claim 37, 38, or 39, wherein R^{604} and R^{605} represent methyl.

41. The compound according to claim 37, 38, or 39, wherein R^{604} represents methyl and R^{605} represents C_{1-4} alkyl substituted by a saturated or unsaturated five- or six-membered carbocyclic or heterocyclic group.

42. The compound according to claim 36, wherein X represents N and all of R^{611} , R^{612} , R^{613} , R^{614} , and R^{615} represent a hydrogen atom, or any one of R^{611} , R^{612} , R^{613} , R^{614} , and R^{615} represents a group other than a hydrogen atom and the remaining groups represent a hydrogen atom.


43. The compound according to claim 42, wherein all of R^{611} , R^{612} , R^{613} , R^{614} , and R^{615} represent a hydrogen atom, or any one of R^{611} , R^{612} , R^{613} , R^{614} , and R^{615} represents C_{1-6} alkyl, $-OR^{616}$, a halogen atom, or nitro and the remaining groups represent a hydrogen atom.

44. The compound according to claim 43, wherein R^{611} represents methoxy and R^{612} , R^{613} , R^{614} , and R^{615} represent a hydrogen atom, or R^{612} represents a bromine atom or methoxy and R^{611} , R^{613} , R^{614} , and R^{615} represent a hydrogen atom, or R^{613} represents a bromine atom, a chlorine atom, a fluorine atom, methyl, methoxy, or nitro and R^{611} , R^{612} , R^{614} , and R^{615} represent a hydrogen atom.

45. The compound according to claim 42, 43, or 44, wherein R^{604} and R^{605} represent methyl.

46. The compound according to claim 42, 43, or 44, wherein R^{604} represents methyl and R^{605} represents C_{1-4} alkyl substituted by a saturated or unsaturated five- or six-membered carbocyclic or heterocyclic group.

47. The compound according to claim 1, represented by formula (If)



wherein

X represents CH or N,

R^{701} represents a hydrogen atom, a fluorine atom at 2-position, a fluorine atom at 3-position, methoxy at 2-position, methoxy at 3-position, or methyl at 2- and 5-positions,

R^{702} represents C_{1-4} alkyl,

R^{703} represents imidazolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, or isothiazolyl in which one or two hydrogen atoms on the groups are optionally substituted by a halogen atom or C_{1-4} alkyl, and

R^{704} and R^{705} , which may be the same or different, represent a hydrogen atom; hydroxyl; nitro; cyano; a halogen atom; $-NR^{706}R^{707}$ wherein R^{706} and R^{707} , which may be the same or different, represent a hydrogen atom or C_{1-4} alkyl in which the alkyl group is optionally substituted by hydroxyl, $-OR^{708}$

wherein R^{708} represents C_{1-4} alkyl, or $-NR^{709}R^{710}$ wherein R^{709} and R^{710} , which may be the same or different, represent a hydrogen atom or C_{1-4} alkyl; $-CONR^{711}R^{712}$ wherein R^{711} and R^{712} , which may be the same or different,

represent a hydrogen atom or C_{1-4} alkyl in which the alkyl group is optionally substituted by hydroxyl, $-OR^{713}$ wherein R^{713} represents C_{1-4} alkyl, or $-NR^{714}R^{715}$ wherein R^{714} and R^{715} , which may be the same or different,


represent a hydrogen atom or C_{1-4} alkyl; $-COOR^{716}$ wherein R^{716} represents a hydrogen atom or C_{1-4} alkyl in which the alkyl group is optionally substituted by hydroxyl, $-OR^{717}$ wherein R^{717} represents C_{1-4} alkyl, or $-NR^{718}R^{719}$

wherein R^{718} and R^{719} , which may be the same or different, represent a hydrogen atom or C_{1-4} alkyl; C_{1-6} alkyl; C_{2-6} alkenyl; C_{2-6} alkynyl; or C_{1-6} alkoxy, in which the alkyl, alkenyl, alkynyl, and alkoxy groups are optionally

substituted by hydroxyl, a halogen atom, $-OR^{720}$ in which R^{720} represents C_{1-4} alkyl, $-NR^{721}R^{722}$ wherein R^{721} and R^{722} , which may be the same or different, represent a hydrogen atom or C_{1-4} alkyl in which the alkyl group is

optionally substituted by hydroxyl or $-OR^{723}$ wherein R^{723} represents C_{1-4} alkyl, or a saturated or unsaturated three- to seven-membered carbocyclic or heterocyclic group in which the carbocyclic and heterocyclic groups are optionally substituted by one or two halogen atoms or C_{1-4} alkyl.

48. The compound according to claim 47, wherein X represents CH, and R^{702} represents




49. The compound according to claim 48, wherein R⁷⁰² represents methyl.

50. The compound according to claim 48 or 49, wherein R⁷⁰⁴ and R⁷⁰⁵ represent methoxy.

51. The compound according to claim 48 or 49, wherein R⁷⁰⁴ represents methoxy, and R⁷⁰⁵ represents C₁₋₄ alkoxy substituted by a saturated or unsaturated five- or six-membered carbocyclic or heterocyclic group.

52. The compound according to claim 47, wherein X represents CH, and R⁷⁰² represents




53. The compound according to claim 52, wherein R⁷⁰² represents methyl.

54. The compound according to claim 52 or 53, wherein R⁷⁰⁴ and R⁷⁰⁵ represent methoxy.

55. The compound according to claim 52 or 53, wherein R⁷⁰⁴ represents methoxy, and R⁷⁰⁵ represents C₁₋₄ alkoxy substituted by a saturated or unsaturated five- or six-membered carbocyclic or heterocyclic group.

56. The compound according to claim 47, wherein X represents N, and R⁷⁰² represents




57. The compound according to claim 56, wherein R⁷⁰² represents methyl.


58. The compound according to claim 56 or 57, wherein R⁷⁰⁴ and R⁷⁰⁵ represent methoxy.

59. The compound according to claim 56 or 57, wherein R⁷⁰⁴ represents methoxy, R⁷⁰⁵ represents C₁₋₄ alkoxy substituted by a saturated or unsaturated five- or six-membered carbocyclic or heterocyclic group.

60. The compound according to claim 47, wherein X represents N, and R⁷⁰² represents



61. The compound according to claim 60, wherein R^{702} represents methyl.
62. The compound according to claim 60 or 61, wherein R^{704} and R^{705} represent methoxy.
63. The compound according to claim 60 or 61, wherein R^{704} represents methoxy, and R^{705} represents C_{1-4} alkoxy substituted by a saturated or unsaturated five- or six-membered carbocyclic or heterocyclic group.
64. The compound according to claim 1, represented by formula (Ig)



wherein

X represents CH or N,

R^{801} represents a hydrogen atom, a fluorine atom at 2-position, a fluorine atom at 3-position, a chlorine atom at 2-position, a chlorine atom at 3-position, methyl at 2- and 3-positions, methyl at 2- and 5-positions, methoxy at 2-position, methoxy at 3-position, methyl at 2-position, or trifluoromethyl at 2-position,

R^{802} represents C_{1-4} alkyl,

R^{804} and R^{805} , which may be the same or different, represent a hydrogen atom; hydroxyl; nitro; cyano; a halogen atom; $-NR^{806}R^{807}$ wherein R^{806} and R^{807} , which may be the same or different, represent a hydrogen atom or C_{1-4} alkyl in which the alkyl group is optionally substituted by hydroxyl, $-OR^{808}$ wherein R^{808} represents C_{1-4} alkyl, or $-NR^{809}R^{810}$ wherein R^{809} and R^{810} , which may be the same or different, represent a hydrogen atom or C_{1-4} alkyl; $-CONR^{811}R^{812}$ wherein R^{811} and R^{812} , which may be the same or different, represent a hydrogen atom or C_{1-4} alkyl in which the alkyl group is optionally substituted by hydroxyl, $-OR^{813}$ wherein R^{813} represents C_{1-4} alkyl, or $-NR^{814}R^{815}$ wherein R^{814} and R^{815} , which may be the same or different, represent a hydrogen atom or C_{1-4} alkyl; $-COOR^{816}$ wherein R^{816} represents a hydrogen atom or C_{1-4} alkyl in which the alkyl group is optionally substituted by hydroxyl, $-OR^{817}$ wherein R^{817} represents C_{1-4} alkyl, or $-NR^{818}R^{819}$ wherein R^{818} and R^{819} , which may be the same or different, represent a hydrogen atom or C_{1-4} alkyl; C_{1-6} alkyl; C_{2-6} alkenyl; C_{2-6} alkynyl; or C_{1-6} alkoxy, in which the alkyl, alkenyl, alkynyl, and alkoxy groups are optionally substituted by hydroxyl, a halogen atom, $-OR^{820}$ in which R^{820} represents C_{1-4} alkyl, $-NR^{821}R^{822}$ wherein R^{821} and R^{822} , which may be the same or different, represent a hydrogen atom or C_{1-4} alkyl in which the alkyl group is optionally substituted by hydroxyl or $-OR^{823}$ wherein R^{823} represents C_{1-4} alkyl, or a saturated or unsaturated three- to seven-membered carbocyclic or heterocyclic group in which the carbocyclic and heterocyclic groups are optionally substituted by one or two halogen atoms or C_{1-4} alkyl, and

R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} , which may be the same or different, represent a hydrogen atom; hydroxyl; C_{1-6} alkyl; $-OR^{836}$ wherein R^{836} represents C_{1-4} alkyl; a halogen atom; nitro; or $-NR^{837}R^{838}$ wherein R^{837} and R^{838} , which may be the same or different, represent a hydrogen atom or C_{1-4} alkyl in which the alkyl group is optionally substituted by hydroxyl, $-OR^{839}$ wherein R^{839} represents C_{1-4} alkyl, or $-NR^{840}R^{841}$ wherein R^{840} and R^{841} , which may be the same or different, represent a hydrogen atom or C_{1-4} alkyl.

65. The compound according to claim 64, wherein X represents CH and all of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represent a hydrogen atom, or any one of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represents a group other than a hydrogen atom and the remaining groups represent a hydrogen atom.

66. The compound according to claim 65, wherein all of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represent a hydrogen atom, or any one of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represents C_{1-6} alkyl, $-OR^{836}$, a halogen atom, or nitro and the remaining groups represent a hydrogen atom.
- 5 67. The compound according to claim 65, wherein R^{831} represents methoxy and R^{832} , R^{833} , R^{834} , and R^{835} represent a hydrogen atom, or R^{832} represents a bromine atom or methoxy and R^{831} , R^{833} , R^{834} , and R^{835} represent a hydrogen atom, or R^{833} represents a bromine atom, a chlorine atom, a fluorine atom, methyl, methoxy, or nitro and R^{831} , R^{832} , R^{834} , and R^{835} represent a hydrogen atom.
- 10 68. The compound according to claim 65, 66, or 67, wherein R^{804} and R^{805} represent methoxy.
69. The compound according to claim 65, 66, or 67, wherein R^{804} represents methoxy and R^{805} represents C_{1-4} alkoxy substituted by a saturated or unsaturated five- or six-membered carbocyclic or heterocyclic group.
- 15 70. The compound according to claim 64, wherein X represents CH, R^{802} represents methyl, and all of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represent a hydrogen atom, or any one of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represents a group other than a hydrogen atom and the remaining groups represent a hydrogen atom.
- 20 71. The compound according to claim 70, wherein all of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represent a hydrogen atom, or any one of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represents C_{1-6} alkyl, $-OR^{836}$, a halogen atom, or nitro and the remaining groups represent a hydrogen atom.
- 25 72. The compound according to claim 70, wherein R^{831} represents methoxy and R^{832} , R^{833} , R^{834} , and R^{835} represent a hydrogen atom, or R^{832} represents a bromine atom or methoxy and R^{831} , R^{833} , R^{834} , and R^{835} represent a hydrogen atom, or R^{833} represents a bromine atom, a chlorine atom, a fluorine atom, methyl, methoxy, or nitro and R^{831} , R^{832} , R^{834} , and R^{835} represent a hydrogen atom.
73. The compound according to claim 70, 71, or 72, wherein R^{804} and R^{805} represent methoxy.
- 30 74. The compound according to claim 70, 71, or 72, wherein R^{804} represents methoxy and R^{805} represents C_{1-4} alkoxy substituted by a saturated or unsaturated five- or six-membered carbocyclic or heterocyclic group.
75. The compound according to claim 64, wherein X represents N and all of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represent a hydrogen atom, or any one of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represents a group other than a hydrogen atom and the remaining groups represent a hydrogen atom.
- 35 76. The compound according to claim 75, wherein all of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represent a hydrogen atom, or any one of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represents C_{1-6} alkyl, $-OR^{836}$, a halogen atom, or nitro and the remaining groups represent a hydrogen atom.
- 40 77. The compound according to claim 75, wherein R^{831} represents methoxy and R^{832} , R^{833} , R^{834} , and R^{835} represent a hydrogen atom, or R^{832} represents a bromine atom or methoxy and R^{831} , R^{833} , R^{834} , and R^{835} represent a hydrogen atom, or R^{833} represents a bromine atom, a chlorine atom, a fluorine atom, methyl, methoxy, or nitro and R^{831} , R^{832} , R^{834} , and R^{835} represent a hydrogen atom.
- 45 78. The compound according to claim 75, 76, or 77, wherein R^{804} and R^{805} represent methoxy.
79. The compound according to claim 75, 76, or 77, wherein R^{804} represents methoxy and R^{805} represents C_{1-4} alkoxy substituted by a saturated or unsaturated five- or six-membered carbocyclic or heterocyclic group.
- 50 80. The compound according to claim 64, wherein X represents N, R^{802} represents methyl, and all of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represent a hydrogen atom, or any one of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represents a group other than a hydrogen atom and the remaining groups represent a hydrogen atom.
- 55 81. The compound according to claim 80, wherein all of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represent a hydrogen atom, or any one of R^{831} , R^{832} , R^{833} , R^{834} , and R^{835} represents C_{1-6} alkyl, $-OR^{836}$, a halogen atom, or nitro and the remaining groups represent a hydrogen atom.

82. The compound according to claim 80, wherein R⁸³¹ represents methoxy and R⁸³², R⁸³³, R⁸³⁴, and R⁸³⁵ represent a hydrogen atom, or R⁸³² represents a bromine atom or methoxy and R⁸³¹, R⁸³³, R⁸³⁴, and R⁸³⁵ represent a hydrogen atom, or R⁸³³ represents a bromine atom, a chlorine atom, a fluorine atom, methyl, methoxy, or nitro and R⁸³¹, R⁸³², R⁸³⁴, and R⁸³⁵ represent a hydrogen atom.

83. The compound according to claim 80, 81, or 82, wherein R⁸⁰⁴ and R⁸⁰⁵ represent methoxy.

84. The compound according to claim 80, 81, or 82, wherein R⁸⁰⁴ represents methoxy and R⁸⁰⁵ represents C₁₋₄ alkoxy substituted by a saturated or unsaturated five- or six-membered carbocyclic or heterocyclic group.

85. The compound according to claim 1, which is a compound selected from a group of the following compounds, or a pharmaceutically acceptable salt or solvate thereof:

- (17) N-{4-[(6,7-dimethoxy-4-quinolyl)oxy]-2-methoxyphenyl}-N'-[(1S)-1-(4-fluorophenyl)ethyl]urea;
- (74) N-{4-[(6,7-dimethoxy-4-quinolyl)oxy]-2-methoxyphenyl}-N'-[1-(1,3-thiazol-2-yl)ethyl]urea;
- (75) N-{4-[(6,7-dimethoxy-4-quinolyl)oxy]-2-methoxyphenyl}-N'-[(1S)-1-(1,3-thiazol-2-yl)ethyl]urea; and
- (76) N-{4-[(6,7-dimethoxy-4-quinolyl)oxy]-2-methoxyphenyl}-N'-[(1R)-1-(1,3-thiazol-2-yl)ethyl]urea.

86. A pharmaceutical composition comprising a compound according to any one of claims 1 to 85 or a pharmaceutically acceptable salt or solvate thereof as an active ingredient.

87. The pharmaceutical composition according to claim 86, which is used in the treatment and prevention of a disease for which the inhibition of macrophage colony-stimulating factor receptor autophosphorylation is effective therapeutically.

88. The pharmaceutical composition according to claim 87, wherein the disease for which the inhibition of macrophage colony-stimulating factor receptor autophosphorylation is effective therapeutically is bone metastasis of malignant tumors including breast cancer, prostatic cancer, and lung cancer; multiple myeloma; osteoporosis; Behcet's disease; or rheumatoid arthritis.

89. Use of a compound according to any one of claims 1 to 85 or a pharmaceutically acceptable salt or solvate thereof, for the manufacture of an agent used in the treatment and prevention of a disease for which the inhibition of macrophage colony-stimulating factor receptor autophosphorylation is effective therapeutically.

90. Use according to claim 89, wherein the disease for which the inhibition of macrophage colony-stimulating factor receptor autophosphorylation is effective therapeutically is bone metastasis of malignant tumors including breast cancer, prostatic cancer, and lung cancer; multiple myeloma; osteoporosis; Behcet's disease; or rheumatoid arthritis.

91. A method for treating and preventing a disease for which the inhibition of macrophage colony-stimulating factor receptor autophosphorylation is effective therapeutically, said method comprising the step of administering a therapeutically or prophylactically effective amount of a compound according to any one of claims 1 to 85 or a pharmaceutically acceptable salt or solvate thereof to a mammal.

92. The method for treating and preventing according to claim 91, wherein the disease for which the inhibition of macrophage colony-stimulating factor receptor autophosphorylation is effective therapeutically is bone metastasis of malignant tumors including breast cancer, prostatic cancer, and lung cancer; multiple myeloma; osteoporosis; Behcet's disease; or rheumatoid arthritis.




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP03/05593

A. CLASSIFICATION OF SUBJECT MATTER Int.Cl. ⁷ C07D215/22, 417/12, A61K31/47, 31/4709, 31/5377, A61P19/02, 19/10, 29/00, 35/00, 35/04, 43/00 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) Int.Cl. ⁷ C07D215/22, 417/12, A61K31/47, 31/4709, 31/5377, A61P19/02, 19/10, 29/00, 35/00, 35/04, 43/00 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) REGISTRY (STN), CAPLUS (STN), CAOLD (STN)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 860433 A1 (KIRIN BEER KABUSHIKI KAISHA), 26 August, 1998 (26.08.98), Full text; particularly, compound 231 & WO 97/17329 A1 & AU 9673400 A & US 6143764 A	1-90
X	EP 1153920 A1 (KIRIN BEER KABUSHIKI KAISHA), 14 November, 2001 (14.11.01), Full text & WO 00/43366 A1 & CA 2361057 A & BR 2000007656 A & NO 2001002617 A	1-90
X	WO 02/32872 A1 (Eisai Co., Ltd.), 25 April, 2002 (25.04.02), Full text & AU 2001095986 A	1-90
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed		"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family
Date of the actual completion of the international search 15 July, 2003 (15.07.03)		Date of mailing of the international search report 29 July, 2003 (29.07.03)
Name and mailing address of the ISA/ Japanese Patent Office Facsimile No.		Authorized officer Telephone No.

Form PCT/ISA/210 (second sheet) (July 1998)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP03/05593

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	WO 03/033472 A1 (KIRIN BEER KABUSHIKI KAISHA), 24 April, 2003 (24.04.03), Full text (Family: none)	1-90

Form PCT/ISA/210 (continuation of second sheet) (July 1998)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP03/05593

Box I Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

- 1.
- ☒
- Claims Nos.: 91, 92

because they relate to subject matter not required to be searched by this Authority, namely:

Claims 91, 92 pertain to methods for treatment of the human body by surgery or therapy and diagnostic methods and thus relates to a subject matter which this International Searching Authority is not required to search.

- 2.
- ☐
- Claims Nos.:

because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

- 3.
- ☐
- Claims Nos.:

because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest ☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.